PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶:
A01N 63/00, 43/04, C12N 15/00, C07H
21/02

(11) International Publication Number:

WO 99/26480

(43) International Publication Date:

3 June 1999 (03.06,99)

(21) International Application Number:

PCT/US98/24950

A1

(22) International Filing Date:

20 November 1998 (20.11.98)

(30) Priority Data:

08/975,424

20 November 1997 (20.11.97) US

(71) Applicants: GENETIX PHARMACEUTICALS, INC. [US/US]; 840 Memorial Drive, Cambridge, MA 02139 (US). MASSACHUSETTS INSTITUTE OF TECHNOLOGY [US/US]; 77 Massachusetts Avenue, Cambridge, MA 02139 (US).

(72) Inventors: LEBOULCH, Philippe; Flagship Warf, Unit 729, 197 8th Street, Charlestown, MA 02129 (US). PAWLIUK, Robert, James; Apartment 3, 52 Maple Avenue, Cambridge, MA 02129 (US). BACHELOT, Thomas; 53, rue Pierre Brunier, F-69300 Caluire (FR).

(74) Agent: CLARK, Paul, T.; Clark & Elbing LLP, 176 Federal Street, Boston, MA 02110-2214 (US).

(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, HR, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published

With international search report.

(54) Title: ANTI-ANGIOGENIC GENE THERAPY VECTORS AND THEIR USE IN TREATING ANGIOGENESIS-RELATED DISEASES

(57) Abstract

A method for inhibiting tumor growth in a human patient harboring a solid tumor, said method comprising administering to said patient a nucleic acid molecule which expresses in said patient an anti-angiogenic polypeptide selected from the group consisting of human angiostatin, murine angiostatin, human endostatin, murine endostatin, and angiogenesis-inhibiting fragments thereof, wherein expression of the anti-angiogenic polypeptide in the patient inhibits angiogenesis in the vicinity of the tumor and/or systemically by diffusion of the recombinant protein to the vascular compartment from secreting transduced cells, thereby inhibiting its growth.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
ΑU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
ΑZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav	TM	Turkmenistan
BF	Burkina Faso	GR	Greece .		Republic of Macedonia	TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
ВJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Ītaly	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG.	Kyrgyzstan	NO. ~	Norway	z₩	Zimbatfwe
CI	Côte d'Ivoire	KP	Democratic People's	NZ	New Zealand		
CM	Cameroon		Republic of Korea	PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

WO 99/26480 PCT/US98/24950

ANTI-ANGIOGENIC GENE THERAPY VECTORS AND THEIR USE IN TREATING ANGIOGENESIS-RELATED DISEASES

Field of the Invention

This invention relates generally to gene therapy for, e.g., cancer.

Background of the Invention

Angiogenesis is the process by which new capillaries are formed from existing vasculature. It is a complex process which involves proliferation and migration of endothelial cells. It plays a fundamental role in reproduction, development and wound repair. Unregulated angiogenesis, however, can further the progression of many diseases, including tumor growth and metastasis, arthritis, diabetes, and some forms of blindness. For example, there is experimental evidence that limits of tumor size and growth are not the failure of the tumor cells to proliferate, but rather a failure of the tumor to provide sufficient nutrients and waste removal to its constituent cells by recruiting surrounding vasculature.

Summary of the Invention

The invention features a method for inhibiting tumor growth in a

human patient harboring a solid tumor, involving administering to the patient a
nucleic acid molecule which expresses in the patient an anti-angiogenic
polypeptide selected from the group consisting of human angiostatin, murine
angiostatin, human endostatin, murine endostatin, and angiogenesis-inhibiting
fragments thereof, wherein expression of the anti-angiogenic polypeptide in the
patient inhibits angiogenesis in the vicinity of the tumor and/or systemically by
diffusion of the recombinant protein to the vascular compartment from
secreting transduced cells, thereby inhibiting its growth.

In a second, related aspect, the invention features tumor inhibition, of

5

10

WO 99/26480 PCT/US98/24950

-2-

the type just described, using nucleic acids molecules of the formula A-B, where A and B are polypeptide and/or export signal joined by a peptide bond; peptide A contains at least 100 amino acids and includes at least kringles 1, 2, and 3 of human or murine angiostatin; and peptide B contains at least 100 amino acids and includes at least 75% of the amino acid sequence of human or murine endostatin. Expression of the fusion anti-angiogenic polypeptide in the patient inhibits angiogenesis in the vicinity of the tumor and/or systemically by diffusion of the recombinant protein to the vascular compartment from secreting transduced cells, thereby inhibiting its growth. In some embodiments of this hybrid polypeptide and/or export signal method, polypeptide and/or export signal A further includes kringle region 4 of angiostatin, and can also include kringle region 5 of plasminogen (the larger protein molecule of which angiostatin is a portion).

In both aspects of the invention, the nucleic acid molecule preferably constitutes a portion of a viral vector or a plasmid, which can either be administered to the patient so that cells of the patient in the vicinity of the tumor and/or systemically by diffusion of the recombinant protein to the vascular compartment from secreting transduced cells are infected or transfected with the nucleic acid encoding the angiogenesis-inhibiting polypeptide, or cells (of the patient, or another human donor, or an animal) are infected or transfected *ex-vivo*, and those infected or transfected cells are then infused into the patient so that the anti-angiogenic polypeptide is expressed in the vicinity of the tumor and/or systemically by diffusion of the recombinant protein to the vascular compartment from secreting transduced cells.

As will be discussed in more detail below, in particularly effective embodiments, the nucleic acid molecule includes a nucleotide sequence

5

10

15

20

PCT/US98/24950

5

10

encoding a preactivation polypeptide and/or export signal for effecting Golgi and/or endoplasmic reticulum export of the anti-angiogenic polypeptide.

In another aspect, the invention features a method for treating a human patient suffering from diabetic retinopathy, involving administering to the patient one of the nucleic acid molecules described above.

The above and other features, objects and advantages of the present invention will be better understood by a reading of the following specification in conjunction with the drawings.

Brief Description of the Drawings

Fig. 1 depicts the structural relationship of angiostatin with plasminogen.

Fig. 2 depicts the structural relationship of endostatin with collagen type XVIII.

Fig. 3 depicts various viral (A. MSCV murine retrovirus; B. Adenoassociated virus; C. HIV based retrovirus; E. recombinant adeno-virus) and
non-viral (D. plasmid) vectors used in the construction of gene therapy vectors
for this invention.

References:

A. Hawley, R.G., F. H. L. Lieu, A. Z. C. Fong, and T. S. Hawley. 1994. Versatile retroviral vectors for potenital use in gene therapy. Gene Therapy 1:136.

B. Hargrove, P. H., E. F. Vanin, G. J. Kurtzman and A. W. Nienhuis. 1997. High-level globin gene expression mediated by a recombinant adeno-associated virus genome that contains a 3' γ globin gene regulatory element and integrates as tandem copies in erythroid cells. Blood 89:2167.

C. Naldini, L., U. Blomer, P. Gallay, D. Ory, R. Mulligan, F. H. Gage, I. M. Verma, and D. Trono. 1996. In vivo gene delivery and stable transduction of nondividing cells by a Lentiviral vector. Science 272: 263.

E. Ohashi, T., K. Watabe, K. Uehara, W. S. Sly, C. Vogler, and Y. Eto. 1997. Adenovirus-mediated gene transfer and expression of human β-glucuronidase gene in the liver, spleen, and central nervous system in mucopolysaccharide type VII mice. PNAS 94: 1287.

MSCV: Murine Stem Cell Virus
LTR: Long Terminal Repeat
RSV: Rous Sarcoma Virus
ITR: Inverted Terminal Repeat

HIV: Human Immunodeficiency Virus IRES: Internal Ribosomal Entry Site GFP: Green Fluorescence Protein HBPRE: Hepatitis B Export Element RRE: Rev Response Element polyadenylation site

w+: yiral packaging sequence

The inverted triangle shows the site at which the anti-angiogenic constructs will be inserted using engineered MluI and XhoI restriction sites.

* denotes specific mutations within the long terminal repeat and leader which bestows the ability for expression in embryonic stem and hematopoietic stem cells.

The arrow denotes the direction of transcription.

Fig. 4 depicts in the left (A) panel nude mice which were implanted with human neuroblastoma cells (line SK-N-AS) transduced with a mock virus and in the right (B) panel, nude mice which were transplanted with human neuroblastoma cells transduced with a retroviral gene therapy vector encoding an angiostatin-endostatin fusion protein.

Fig. 5 shows the nucleotide sequence (SEQ ID NO: 1) and amino

10

15

20

25

acid sequence (SEQ ID NO: 2) of human plasminogen and the nucleotide sequence (SEQ ID NO: 5) and amino acid sequence (SEQ ID NO: 6) of human angiostatin.

Fig. 6 shows the nucleotide sequence (SEQ ID NO: 9) and amino acid sequence (SEQ ID NO: 10) of murine endostatin.

Fig. 7 shows the nucleotide sequence (SEQ ID NO: 3) and amino acid sequence (SEQ ID NO: 4) of murine plasminogen and the nucleotide (SEQ ID NO: 7) and amino acid sequence (SEQ ID NO: 8) of murine angiostatin.

Detailed Description

This invention provides gene therapy using a vector having a nucleotide sequence encoding one of the above-identified anti-angiogenic polypeptides. Described below in more detail are some of the components of the vectors and methods of the invention.

By a gene therapy vector is meant a vector useful for gene therapy. Gene therapy vectors carry a gene of interest that is useful for gene therapy. The gene therapy vectors are able to be transferred to the cells of an animal, e.g., a human, and are able to express the gene of interest in such cells so as to effect gene therapy. The vector can be, e.g., chromosomal, non-chromosomal, or synthetic, and can be RNA or DNA. The vector can be, e.g., a plasmid, a virus or a phage. Preferred vectors include, e.g., retroviral vectors, adenoviral vectors, adeno-associated vectors, herpes virus vectors, Simliki Forest Virus-based vector, Human Immunodeficiency virus, Simian Immunodeficiency virus, and non-viral plasmids. A preferred retroviral vector is Murine Stem Cell Virus (MSCV), which is a variant of Moloney Murine Leukemia Virus (MoMLV).

By anti-angiogenic polypeptide is meant a polypeptide which inhibits

10

15

20

25

angiogenesis. The terms polypeptide, protein and polypeptide and/or export signal are used interchangeably herein. By angiogenesis is meant the process by which new vasculature, in particular, new capillaries, are formed from existing vasculature. Angiogenesis is a complex process entailing numerous steps, including local dissolution of the basement membrane, migration of endothelial cells into the surrounding stroma, proliferation of the endothelial cells at the leading edge to form a migrating column of cells, branching and fusion of the newly formed vascular loops, and formation of a new basement membrane. By inhibiting angiogenesis is meant completely or partially inhibiting the formation of such new vasculature.

In certain embodiments, the anti-angiogenic polypeptide is an antiangiogenic fragment of plasminogen (in particular, angiostatin), an antiangiogenic fragment of collagen XVIII (endostatin) or a fusion of the two fragments.

Angiostatin is an internal fragment of plasminogen having a molecular weight of 38 or 45 kDa, depending on whether it contains kringles 1-3 or 1-4. In the invention, either can be used, or a molecule including kringles 1-3 and a portion of kringle 4 can be used. Angiostatin can be naturally produced in vivo in small amounts by tumor cells, e.g. murine Lewis lung carcinoma cells, by proteolytic cleavage of plasminogen so as to eliminate the N-terminal portion including the signal polypeptide and/or export signal and the preactivation polypeptide and/or export signal, as well as the C-terminal portion following kringle 3 or 4. Mouse and human angiostatin have been purified and sequenced. In preferred embodiments, the gene therapy vectors of this invention encode angiostatin having kringles 1, 2, 3 and 4.

In another preferred embodiment, the anti-angiogenic polypeptide is

WO 99/26480 PCT/US98/24950

-7-

endostatin or a biologically active analog or fragment thereof. Endostatin can be naturally produced *in vivo* in small amounts by tumor cells, e.g., murine angiosarcoma cells, by proteolytic cleavage of endogenous collagen XVIII so as to eliminate the N-terminal portion including the signal polypeptide and/or export signal and the preactivation polypeptide and/or export signal, as well as the C-terminal portion following kringle 3 or 4. See Fig.2. Mouse endostatin has been sequenced, and the human molecule (SEQ ID NOs: 17 and 18) forms a portion of collagen 18 (SEQ ID NOs: 19 and 20).

The human molecule position and sequence are apparent from an alignment of the active, Lys-terminated active region of human collagen 18 with murine endostatin, such that the C-terminal lysine residues align, bringing the active endostatin sequences into alignment.

In yet another preferred embodiment, the anti-angiogenic polypeptide is an in-frame fusion of angiostatin or a biologically active analog or fragment thereof and endostatin or a biologically active analog or fragment thereof. Preferably, the angiostatin or biologically active analog or fragment is 5' of the endostatin or biologically active analog or fragment. In certain embodiments, the angiostatin-endostatin fusion proteins exhibit synergistic anti-angiogenic properties.

By fragment is meant some portion of the naturally occurring antiangiogenic polypeptide. Preferably, the fragment is at least 20 amino acid residues, more preferably at least 50 amino acid residues, and most preferably at least 100 amino acid residues in length. Fragments include chimeric constructs composed of at least a portion of the relevant gene and another molecule. The ability of a candidate fragment to exhibit a biological activity of the anti-angiogenic polypeptide can be assessed by methods known to those skilled in the art, e.g., by its ability to inhibit proliferation of bovine capillary

5

10

15

20

10

15

20

25

cells, or by its ability to inhibit growth of primary tumor cells, e.g., as described herein. See, e.g., Example 9. Also included are fragments containing residues that are not required for biological activity of the fragment or that result from alternative mRNA splicing or alternative protein processing events.

Internal or terminal fragments of a polypeptide can be generated by removing one or more nucleotides from one end (for a terminal fragment) or both ends (for an internal fragment) of a nucleic acid which encodes the polypeptide.

In preferred embodiments, the gene therapy vector of this invention is capable of hybridizing to the native anti-angiogenesis polypeptide-encoding regions and has at least about 80%, preferably at least about 90%, and more preferably at least about 95%, sequence identity to the native nucleotide sequences, and encodes a polypeptide which has anti-angiogenic activity; or a biologically active fragment of any of the above nucleotide sequences wherein the encoded polypeptide has anti-angiogenic activity.

The nucleotide sequences of the present invention can be in the form of RNA or DNA, and the nucleotide sequence can be double-stranded or single stranded and, if single stranded, can be the coding strand or non-coding (antisense) strand.

The coding sequence which encodes the anti-angiogenic polypeptide can be identical to the native coding sequences, or can be a different coding sequence which, as a result of the degeneracy of the genetic code, encodes the same anti-angiogenic polypeptide.

In certain embodiments, the gene therapy vector also has a nucleotide sequence encoding a signal polypeptide and/or export signal (SP) for effecting secretion of the anti-angiogenic polypeptide. Examples of signal polypeptide and/or export signal include plasminogen signal polypeptide and/or export

10

15

20

25

signal. Preferably, the signal polypeptide and/or export signal is 5' (i.e., upstream) of the nucleotide sequence encoding the anti-angiogenic polypeptide.

Preferably, the gene therapy vector has a nucleotide sequence encoding a preactivation polypeptide and/or export signal (PAP), which is a small polypeptide and/or export signal which effects folding and secretion of the anti-angiogenic polypeptide *in vivo*. Examples of preactivation polypeptide and/or export signal include plasminogen preactivation polypeptide and/or export signal, described herein, and PAP's of other proteins in the blood clotting cascade.

Preferably, the preactivation polypeptide and/or export signal is positioned 5' of the nucleotide sequence encoding the anti-angiogenic polypeptide. In embodiments which have a signal sequence and an anti-angiogenic polypeptide, preferably the preactivation polypeptide and/or export signal is 5' of the nucleotide sequence encoding the anti-angiogenic polypeptide, and 3' of the nucleotide sequence encoding the signal polypeptide and/or export signal.

We have discovered that results obtained using constructs containing a PAP- encoding nucleic acid sequence are far superior to results using constructs lacking a PAP-encoding sequence. Our hypothesis to explain these unexpectedly superior results with PAP is that, during the complex process by which the anti-angiogenic polypeptide is expressed and processed in living cells, the PAP polypeptide and/or export signal facilitates the export of the polypeptide from the cellular Golgi apparatus and/or the endoplasmic reticulum (ER). The corollary is that, absent PAP, a significant portion of the expressed polypeptide remains trapped in the Golgi and/or ER.

The PAP exemplified herein is derived from human plasminogen; this

-10-

PAP is currently preferred. Our discovery that the use of a PAP dramatically improves results leads us to believe that other PAP's would be useful as well, and such others are therefore contemplated for use in the invention. Thus, as used herein, "PAP" refers to a polypeptide and/or export signal which is naturally associated with a eukaryotic (preferably human) protein, the exportation of which is facilitated by its associated PAP. Examples of other human proteins whose Golgi/ER export is PAP-facilitated include other secreted proteins of the blood coagulation cascade, e.g., fibrinogen, prothrombin, Factor VIII, and Factor IX. Other secreted human proteins also are associated with potentially useful PAPs.

It is not essential that the PAP used in the invention be identical in amino acid sequences to a native PAP; it is well-known that polypeptide and/or export signal that facilitate protein secretion or export, e.g., signal polypeptide and/or export signal and PAPs, can vary from the native forms to a certain extent and still retain their function. Therefore, PAPs useful according to the invention preferably have 75% or greater amino acid sequence identity with a native PAP.

In certain embodiments, the gene therapy vector has a nucleotide sequence encoding a tag for identification of the anti-angiogenic polypeptide and/or export signal. In certain embodiments, the tag is 5' of the nucleotide sequence encoding the anti-angiogenic polypeptide; in other embodiments, the tag is 3' of the nucleotide sequence encoding the anti-angiogenic polypeptide. In embodiments in which the anti-angiogenic polypeptide is endostatin or an angiostatin-endostatin fusion, it is preferred that the tag be 5' of the nucleotide sequence encoding endostatin.

In certain embodiments the gene therapy vector includes a selectable

5

10

15

20

marker, e.g., a Neomycin phosphotransferase gene, or a humanized red-shifted green fluorescent protein.

The invention also includes a cell infected or transfected with a gene therapy vector described herein. Preferably, the cell is an animal cell, more preferably an autologous or allogeneic human cell. The gene therapy vectors described herein can be introduced into a cell, e.g., by transformation, transfection, transduction, infection, or ex vivo injection. They can be targeted to a particular cell type.

Administration of nucleic acid, e.g., a gene therapy vector, can be accomplished by any method which allows the nucleic acid to reach the target cells. These methods include, e.g., injection, deposition, implantation, suppositories, oral ingestion, inhalation, topical administration, or any other method of administration where access to the target cells by the nucleic acid is achieved. Injections can be, e.g., intravenous, intradermal, subcutaneous, intramuscular or intraperitoneal. Implantation includes inserting implantable drug delivery systems, e.g., microspheres, hydrogels, polymeric reservoirs, cholesterol matrices, polymeric systems, e.g., matrix erosion and/or diffusion systems and non-polymeric systems, e.g., compressed, fused or partially fused pellets. Suppositories include glycerin suppositories. Oral ingestion doses can be enterically coated. Inhalation includes administering the nucleic acid with an aerosol in an inhalator, either alone or attached to a carrier that can be absorbed.

In certain embodiments of the invention, administration can be designed so as to result in sequential exposures to the nucleic acid over some time period, e.g., hours, days, weeks, months or years. This can be accomplished by repeated administrations of the nucleic acid, e.g., by one of the methods described above, or alternatively, by a controlled release delivery system in

to a state of the second

5

10

15

20

WO 99/26480 PCT/US98/24950

-12-

which the nucleic acid is delivered to the animal over a prolonged period without repeated administrations. By a controlled release delivery system is meant that total release of the nucleic acid does not occur immediately upon administration, but rather is delayed for some time. Release can occur in bursts or it can occur gradually and continuously. Administration of such a system can be, e.g., by long acting oral dosage forms, bolus injections, transdermal patches or subcutaneous implants. Examples of systems in which release occurs in bursts include, e.g., systems in which the nucleic acid is entrapped in liposomes which are encapsulated in a polymer matrix, the liposomes being sensitive to a specific stimulus, e.g., temperature, pH, light, magnetic field, or a degrading enzyme, and systems in which the nucleic acid agent is encapsulated by an ionically-coated microcapsule with a microcapsule core-degrading enzyme. Examples of systems in which release of the nucleic acid is gradual and continuous include, e.g., erosional systems in which the nucleic acid is contained in a form within a matrix, and diffusional systems in which the nucleic acid permeates at a controlled rate, e.g., through a polymer. Such sustained release systems can be, e.g., in the form of pellets or capsules.

The nucleic acid is administered to the patient in a therapeutically effective amount. By therapeutically effective amount is meant that amount which is capable of at least partially preventing or reversing the disease. A therapeutically effective amount can be determined on an individual basis and will be based, at least in part, on consideration of the patient's size, age, the efficacy of the particular nucleic acid used, the type of delivery system used, the time of administration relative to the onset of disease symptoms, and whether a single, multiple, or controlled release dose regimen is employed. A therapeutically effective amount can be determined by one of ordinary skill in the art employing such factors and using no more than routine experimentation.

5

10

15

20

10

15

20

25

In certain embodiments, a therapeutically effective amount of an antiangiogenic polypeptide is administered by providing to the animal a nucleic acid encoding the polypeptide and expressing the polypeptide in vivo. Nucleic acids encoding the polypeptide, or mutants thereof, can be administered in any biologically effective carrier, e.g. any formulation or composition capable of effectively delivering the nucleotide sequence for the anti-angiogenic polypeptide to cells in vivo. Approaches include, e.g., insertion of the nucleic acid into viral vectors. Viral vectors can be delivered to the cells, e.g., by infection or transduction using the virus. Viral vectors can also be delivered to the cells, e.g., by physical means, e.g., by electroporation, lipids, cationic lipids, liposomes, DNA gun, Ca₃(PO₄)₂ precipitation, or delivery of naked DNA. In certain preferred embodiments, the virus is administered by injection, e.g., intramuscular injection, in a dose range of about 10³ to about 10¹⁰ infectious particles per injection, more preferably in a dose range of about 105 to about 108 infectious particles per injection. Single or multiple doses can be administered over a given period of time, depending, e.g., upon the disease.

An alternative is insertion of the nucleic acid encoding the antiangiogenic polypeptide into a bacterial or eukaryotic plasmid. Plasmid DNA can be delivered to cells with the help of, e.g., cationic liposomes (lipofectinTM; Life Technologies, Inc., Gaithersburg, MD) or derivatized (e.g., antibody conjugated) polylysine conjugates, gramacidin S, artificial viral envelopes or other such intracellular carriers, as well as direct injection of the gene construct or Ca₃(PO₄)₂ precipitation carried out in vivo, or by use of a gene gun. The above-described methods are known to those skilled in the art and can be performed without undue experimentation.

Since transfer of the nucleic acid to appropriate target cells represents the critical first step in gene therapy, choice of the particular gene delivery -14-

system will depend on such factors as the intended target and the route of administration, e.g., locally or systemically. Targets for delivery of the nucleic acid can be, e.g., specific target cells which are diseased. For example, the target can be, e.g., the peritoneal cavity, gastro-intestinal tract, bone marrow cavity, liver, lungs, muscles, vasculature, pericardial cavity, pleural cavity, skin, sub-cutaneous or deep connective tissues, central nervous system, spinal fluid, eye, or specific sites of tumor growth. Administration can be directed to one or more cell types, and to one or more cells within a cell type, so as to be therapeutically effective, by methods known to those skilled in the art. For example, the nucleic acid can be, e.g., coupled to an antibody, to a ligand to a cell surface receptor, or to a toxin component, or can be contained in a particle which is selectively internalized into cells, e.g., liposomes, or a virus where the viral receptor binds specifically to a certain cell type, or a viral particle lacking the viral nucleic acid, or can be administered by local injection.

In certain embodiments, the nucleic acid is administered to the patient by introducing *ex vivo* the nucleic acid into cells of the patient, or into syngeneic or allogeneic or xenogeneic cells, and then administering the cells having the nucleic acid to the animal. Any cell type can be used. In certain embodiments, the cells having the introduced nucleic acid are expanded and/or selected after the nucleic acid transfer. The cells having the transferred nucleic acid are subsequently administered to the patient. Preferably, the cells are administered in a dose range of about 1 x 10⁶ to about 1 x 10⁹ cells/dosage/day, and most preferably at about 1 x 10⁷ to about 1 x 10⁸ cells/dosage/day. The cells can be administered by any method which results in delivering the

25 "transferred nucleic acid in the cells to the desired target. For example, the cells "a can be implanted directly into a specific tissue of the patient, or implanted after encapsulation within an artificial polymer matrix. Examples of sites of

5

10

15

7.4 1.18

5

10

15

20

implantation include, e.g., the peritoneal cavity, gastro-intestinal tract, bone marrow cavity, liver, lungs, muscles, vasculature, pericardial cavity, pleural cavity, skin, sub-cutaneous or deep connective tissues, central nervous system, spinal fluid, eye, or specific sites of tumor growth.

Systemic delivery can be achieved, e.g., by introducing the nucleic acid into cells which circulate in the peripheral blood of the patient, or which give rise to cells which circulate in the peripheral blood. In certain embodiments, the nucleic acid is introduced into such cells ex vivo, and these cells are then administered to the patient, resulting in systemic delivery within the peripheral blood. These cells can be the cells of the patient or allogeneic cells. Preferred cells in which the nucleic acid can be introduced are hematopoietic cells.

In certain embodiments, other therapy is additionally administered. For example, if the animal is being treated for a tumor, other tumor therapy, e.g., another therapeutic agent, chemotherapy, radiation or surgery, is additionally administered to the patient, either simultaneously or at different times.

Treating is meant to include, e.g., preventing, treating, reducing the symptoms of, or curing the disease. I.e. treating a tumor includes preventing growth of the tumor, causing shrinkage of the tumor, or preventing development of micro-metasteses.

Preferably, the recombinant nucleic acid is a gene therapy vector, e.g., as described herein. Preferably, the anti-angiogenic polypeptide is angiostatin, endostatin, an angiostatin-endostatin fusion protein, or biologically active analogs or fragments thereof. In certain embodiments, the angiostatin has kringles 1, 2 and 3; in other embodiments, the angiostatin has kringles 1, 2, 3 25 and 4, and, in some embodiments, kringle 5 of human or murine plasminogen. Angiostatin is described in O'Reilly and Folkman U.S. Patent No. 5,639,725, hereby incorporated by reference. Endostatin is described in O'Reilly and

Folkman PCT Appln. No. WO 97/15666, published May 1, 1997, hereby incorporated by reference.

In certain embodiments, the recombinant nucleic acid has been introduced ex vivo into cells so as to express the anti-angiogenic polypeptide in the cells, and the recombinant nucleic acid is administered to the patient by administering to the patient the cells containing the recombinant nucleic acid. In certain embodiments, the cells are derived from the patient; in other embodiments the cells are allogeneic cells relative to the cells of the patient.

Where cells are infected or transfected *ex vivo* for later infusion into the patient, the cells are preferably hematopoietic cells, but can also be mesenchymal cells, stem cells, epithelial cells (e.g., from the gut), or dendritic cells.

The gene therapy vectors of the invention can be provided in a pharmaceutical composition comprising a therapeutically effective amount of the recombinant nucleic acid together with a pharmaceutically acceptable carrier. Pharmaceutically acceptable carriers include, e.g., water, saline, dextrose, glycerol, ethanol, liposomes and lipid emulsions.

The following non-limiting examples further illustrate the present invention.

20 EXAMPLES

Example 1: Construction of Inserts for Gene Therapy Vectors Containing cDNA for Angiostatin, Endostatin or Angiostatin-Endostatin Fusion Proteins

The following genetic constructs are inserted into retroviral gene therapy.

vectors; the genetic constructs contain human or murine cDNA for angiostatin,
endostatin or an angiostatin-endostatin fusion, and DNA encoding a signal

5

10

-17-

polypeptide and/or export signal (SP), a tag (FLAG), and, preferably, a preactivation polypeptide and/or export signal (PAP). The constructs are all made using standard genetic engineering techniques, and their insertion into retroviral gene therapy vectors is carried out using known methods. The constructs have the following components:

	Murine Constructs	
	SP-K1-K2-K3-Flag	
	SP-K1-K2-K3-K4-Flag	
	SP-K1-K2-K3-K4-K5-Flag	
10	SP-PAP-K1-K2-K3-Flag	
	SP-PAP-K1-K2-K3-K4-Flag	(SEQ ID NO: 11 and 12)
	SP-Flag-Endo	(SEQ ID NO: 13 and 14)
	SP-K1-K2-K3- Flag-Endo	
	SP-K1-K2-K3-K4- Flag-Endo	(SEQ ID NO: 15 and 16)
15	SP-PAP-K1-K2-K3- Flag-Endo	
	Human Constructs	
	Human Constructs SP-K1-K2-K3	
	SP-K1-K2-K3	
20	SP-K1-K2-K3 SP-K1-K2-K3-K4	
20	SP-K1-K2-K3 SP-K1-K2-K3-K4 SP-K1-K2-K3-K4-K5	
20	SP-K1-K2-K3 SP-K1-K2-K3-K4 SP-K1-K2-K3-K4-K5 SP-PAP-K1-K2-K3	
20	SP-K1-K2-K3 SP-K1-K2-K3-K4 SP-K1-K2-K3-K4-K5 SP-PAP-K1-K2-K3 SP-PAP-K1-K2-K3-K4	
20	SP-K1-K2-K3 SP-K1-K2-K3-K4 SP-K1-K2-K3-K4-K5 SP-PAP-K1-K2-K3 SP-PAP-K1-K2-K3-K4 SP-PAP-K1-K2-K3-K4-K5	

-18-

Nucleic acid and amino acid sequences for mouse and human angiostatin and endostatin used in these constructs are shown in Figs. 5-7.

Nucleic acid and amino acid sequence of the FLAG peptide:

amino terminus-5'- ASP TYR LYS ASP ASP ASP LYS GAC TAC AAG GAC GAT GAC AAG

Human plasminogen derivative constructs

The entire coding region of the human plasminogen cDNA from the start (ATG) to

the stop (TAA) codon is 2433bp in size.

This sequence encodes a signal peptide (bp 1-57), a preactivation peptide (bp 58-288), and 5 distinct structural regions known as kringles (K1-K3 from bp 289-1092; K4 from bp1093-1380; K5 from bp 1381-1740). Please note that although I have given precise bp measurements for kringles K4 and K5, it can be argued that the sequence encoding K4 is between bp1056-1440 and the sequence encoding K5 is between bp1362-1680.

A DNA fragment encoding a portion of the human plasminogen protein from bp 1 to 1377 was obtained by PCR of a widely available human liver cDNA library using synthetic DNA oligonucleotides complementary to sequences immediately preceding the signal peptide and immediatly following kringle 4. This fragment contains the sigal peptide (bp1-57), the preactivation peptide (bp 58-288), kringles 1 (bp289-549), 2 (bp 550-804), 3 (bp 805-1092) and 4 (bp 1093-1380). The synthetic oligonucleotides used for this reaction contained engineered recognition sites for the restriction enzymes EcoRI and XhoI. Following the PCR reaction the amplified fragment was cloned into the EcoRI/XhoI sites of BluescriptSK(-) (Stratgene) using standard techniques (Maniatis). Following cloning the integrity of the amplified sequence was verified by sequencing both strands using the Sanger method (Sanger). Various derivatives of the cloned fragment were subsequently constructed using BluescriptSK(-) (Stratagene) as a backbone. A full list of the derivatives are described in Table 1. Briefly, the variations are composed of constructs containing various combinations of kringles with or without the signal and/or preactivation peptide sequences. These derivatives were constructed using both standard techniques as well as PCR and the use of double stranded synthetic oligonucleotides. In all cases the integrity of the start codon, coding sequence and termination codon was verified by double stranded sequencing using the Sanger method.

Murine plasminogen derivative constructs

The coding sequence for murine plasminogen is 2439bp in size and, similar to the human plasminogen cDNA encompasses a sequence encoding signal and preactivation peptides (bp 1-57 and 58-288 respectively) in addition to 5 kringle regions; kringle 1-3 (bp 289-1092), kringle 4 (bp 1093-1380) and kringle 5 (bp 1381-1743). Again, although I have given precise bp measurements for kringles K4 and K5, it can be argued that the sequence encoding K4 is between bp1056-1440 and the sequence encoding K5 is between bp1362- 1680.

The murine plasminogen cDNA has previously been cloned and was made available to us. Derivatives of murine plasminogen were constructed using sequences derived from bp 1-1743 of the coding sequence. Various combinations of kringle regions with or without signal and preactivation peptide regions were made using BluescriptSK(-) (Stratagene, La Jolla, CA) as the vector backbone. These derivatives were constructed using standard cloning techniques (Maniatis, Molecular cloning; a laboratory manual, second edition, 1989) in combination with PCR utilizing synthetic oligonuleotides using

10

15

20

25

Angiostatin function was not altered by adding the FLAG polypeptide and/or export signal to either the N- or C-terminal ends, whereas endostatin was functional only if FLAG was added to its N-terminal end.

Example 2: Construction of Retroviral Gene Therapy Vectors

This example illustrates the construction of retroviral gene therapy vectors comprising cDNA for angiostatin, endostatin or angiostatin-endostatin fusion proteins.

The DNA inserts from Example 1 were inserted into two retroviral vectors. Both vectors were derived from the Murine Stem Cell Virus (MSCV), which is a variant of Moloney Murine Leukemia Virus (MoMLV) having several mutations allowing high, sustained expression in hematopoietic stem cells and their progeny. In both cases, the angiostatin, endostatin, or angiostatin-endostatin fusion DNA inserts were under the transcriptional control of the retroviral left Long Terminal Repeat (LTR). In the first vector, the dominant selectable marker was the Neomycin phosphotransferase gene (NeoR), which confers resistance to G418, and is driven by an internal phosphoglycerate kinase (PGK) promoter. In the second vector, the dominant selectable marker was the humanized, red-shifted green fluorescent protein (EGFP), which is co-translationally expressed by means of an Internal Ribosome Entry Site (IRES) from the Encephalomyocarditis virus (EMCV).

The retroviral gene therapy vectors were transfected by CaPO₄ precipitation in the transient ecotropic packaging cell-line BOSC 23, Pear et al., PNAS 90:8392 (1993). Viral supernatants were collected two days thereafter and filtered through 0.45 mm filters. Filtered viral supernatants were subsequently used to infect GENETIX's stable amphotropic retroviral packaging cell-line AM12 (Genetix Pharmaceuticals, Inc., Cambridge, MA). After another two days, viral supernatants from transduced AM12 were filtered

15

20

25^{rr.}

and used to infect GENETIX's stable ecotropic retroviral packaging cell-line GP+E86 (Genetix Pharmaceuticals, Inc.). Both transduced AM12 and GP+E86 were then selected in the presence of G418 (in the case of constructs bearing NeoR) or sorted by Fluorescent Activated Cell Sorter (FACS) for EGFP expression. Viral titers were estimated according to standard practice by counting G418 resistant colonies among NIH3T3 cells exposed to diluted virus preparation. Ecotropic viral titers were above 5 x 10⁵/ml of viral supernatants, only 3-fold lower than "empty" control vectors. No Replication Competent Retrovirus (RCR) was detected in standard assays.

10 <u>Example 3</u>: <u>Transduction of Target Cells Using Retroviral Gene Therapy Vectors</u>

This example illustrates the stability of retroviral gene therapy vector transmission and the lack of toxicity in non-endothelial target cells.

Following 24-hour incubation of confluent viral producer cells in 100 mm plates, viral supernatant was removed and filtered (0.45 µm filter, Gelman Sciences, Ann Arbor, MI). Viral supernatant, containing 7 µg/ml polybrene (Sigma, St. Louis, MO), was added to target cells 24 hours after plating the target cells. Fresh medium was added after 4-12 hours, and, after an additional 48 hours, cells were selected for retroviral infection by exposure to medium containing 1 mg/ml G418 (Gibco BRL, Grand Island, NY) or by FACS sorting (FACStar cell sorter, Becton Dickinson, San Jose, CA). The stability of transmission of the retroviral gene therapy vectors described in Example 2 was examined by Southern blot analysis of transduced NIH3T3 cells, using specific probes (EGFP) and restriction enzyme digestion of genomic DNA with Sac1, which cuts only once in each LTR. Stable chromosorfial integration of intact proviruses of appropriate length was observed with all constructs.

The lack of non-specific toxicity on non-endothelial cells was

established by using filtered viral supernatants to transduce various tumor cell-types and cell-lines (NIH3T3 cells, K562 cells (ATCC), and human SK-N-AS neuroblastoma cells; Cohen, P.S., *Cancer Research*, <u>55</u>:2380 (1995).

Transduced cell populations were subsequently selected with G418 or sorted for EGFP expression by FACS. No obvious effects on cell viability, growth or other phenotypical characteristics were detected.

Example 4: Protein Expression of Angiostatin, Endostatin and Angiostatin-Endostatin Fusion Proteins in Cells Transduced with Retroviral Gene Therapy Vectors

This example illustrates that recombinant angiostatin, endostatin, and angiostatin-endostatin fusion proteins were readily detected in retrovirally transduced cells and their supernatant, indicating efficient expression and secretion.

MSCV virus based vectors containing sequences encoding murine Kringle 1 (K1), K1K5, K1K2K3, K1K2K3K4, and K1K2K3K4K5 were used to transduce NIH3T3 cells. With regard to the murine recombinant proteins, Western blot analysis of transduced cells and their supernatant was performed by means of a monoclonal antibody that recognizes the FLAG polypeptide and/or export signal. Because this antibody is not mono-specific, significant cross-reactivity with murine proteins was apparent. However, by comparing the pattern obtained with mock cells, it was clear that the antibody revealed an additional band of appropriate size in all transduced cells. Moreover, the recombinant proteins were detected in cell supernatants at levels above 50 ng/ml, using a protein concentration/semi-purification procedure (Centricon columns. Amicon, Beverly, MA), With regard to the human recombinant proteins, no FLAG tag was added, so a monoclonal antibody that recognizes specifically the first three kringles of human plasminogen in its native, non-

10

15

20

25.

10

15

20

-22-

denatured form was used; O'Reilly et al., Cell 79:315 (1994). Because of this constraint, Western blot analysis using denaturing gels could not be performed. An ELISA assay was performed which indicated that human recombinant angiostatin was detected at levels likely to be therapeutic according to previous findings in the model of Lewis Lung Carcinoma Id.

These results indicate that high levels of recombinant proteins of expected length were expressed in retrovirally transduced cells and were efficiently secreted.

Example 5: In Vivo Anti-Tumor Activity of Cells Transduced with Gene
Therapy Vectors Encoding the Angiostatin-Endostatin Fusion
Protein

Human SK-N-AS neuroblastoma cells (Cohen, 1995) were transduced with the retroviral gene therapy vector containing the angiostatin-endostatin fusion protein, described in Example 2. These cells (1,000,000) were suspended in 1 mL Dulbecco's phosphate buffered saline and injected into the right mid-quadrant of nude immuno-compromised mice. While no impairment of the in vitro growth of transduced cells was observed, a dramatic decrease in tumor growth in nude mice cells following subcutaneous implantation of the transduced cells was evident as compared to "mock virus"-transduced control cells.

Example 6: Ex Vivo Transfer of Retroviral Gene Therapy Vectors Encoding
Anti-Angiogenic Polypeptides to Primary Hematopoietic Cells,
and Subsequent Transplantation to Recipient Mice

This example illustrates infection of primary hematopoietic cells from

44 . 2

5

10

15

20

donor mice with retroviral gene therapy vectors encoding angiostatin, endostatin, or an angiostatin-endostatin fusion protein, and a selectable GFP marker, and subsequent transplantation of the transduced hematopoietic cells into recipient mice.

Femoral bone marrow cells are harvested from male donor C57BL6/J-Ly5.1 mice (Jackson Labs, Bar Harbor, ME), intravenously injected four days previously with 150 mg/kg of 5-fluorouracil (5-FU). Bone marrow cells are cultured for two days in medium composed of DMEM, 15% fetal calf serum, 10 ng/ml human IL-6, 6 ng/ml murine IL-3 and 100 ng/ml murine Steel factor prior to two days of culture atop a confluent monolayer of irradiated (1,500 cGy, ¹³⁷Cs γ-irradiation) viral producer cells in the above medium including 6 ug/ml of prolamine sulfate. The viral producer cells are transfected with a retroviral gene therapy vector, as described above. Upon completion of the coculture infection protocol, recovered non-adherent cells are cultured for an additional 48 hours to allow for expression of the transferred GFP gene. Retrovirally transduced cells expressing the transferred GFP gene are subsequently identified and selected for, using a FACStar+ cell sorter (Becton Dickinson, San Jose, CA). The GFP+ cells are intravenously injected into congenic female C57BL6/J-Ly5.2 recipient mice (National Cancer Institute, Washington, DC) previously given 950 cGy (83cGy/min, ¹³⁷Cs γ-rays) of whole body irradiation. In each case, a small fraction of GFP+ sorted cells is used for day 12 CFU-S and in vitro clonogenic progenitor assays to assess the efficiency of the infection and selection procedures on these more mature cell types.

25 Example 7: Engraftment of Recipient Mice with Donor Derived Hematopoietic Cells

This example illustrates engraftment of the recipient mice with the

-24-

donor-derived transfected hematopoietic cells from Example 6.

The donor and recipient mice are phenotypically distinguishable on the basis of Y chromosome specific sequences, as well as on the basis of allelic differences at the murine CD45 cell surface antigen locus. Male donor mice are homozygous for the CD45.2 allele, while female recipient mice are homozygous for CD45.1. The engraftment of recipient mice with donor-derived (CD45.2+) cells is assessed at both short (5 weeks) and long (34 months) time points post-transplant by flow cytometric analysis of peripheral blood samples stained with a phycoerythrin labeled antibody specific for the CD45.2 antigen (Pharmingen, San Diego, CA). The results indicate that engraftment occurs.

Example 8: Proviral Marking and GFP Expression in Recipient Mice

This example illustrates the presence of recombinant provirus and expression of the transferred GFP gene in the recipient mice from Example 6.

The level of proviral marking in reconstituted animals is initially determined by Southern blot and semi-quantitative PCR analysis of DNA obtained from peripheral blood leukocytes. The large majority of donor-derived (CD45.2+) cells in recipient mice contain a minimum of one copy of recombinant provirus. In addition, flow cytometric analysis of peripheral blood leukocytes is performed to ascertain the proportion of cells expressing the transferred GFP cDNA. Because the GFP and angiogenic inhibitor protein cDNAs are both driven from the same regulatory sequences, due to the inclusion of an internal ribosomal entry site (IRES) element, the analysis of GFP expression in the peripheral blood provides an indirect measurement of the levels of anti-angiogenic protein being expressed. The results indicate expression of the transferred genes.

5

10

15

20

10

15

20

25

Example 9: Anti-Angiogenic Polypeptide Expression in Recipient Mice

This example illustrates the presence of anti-angiogenic polypeptide in the sera of the recipient mice from Example 6, using both physical and functional assays.

Serum obtained from the transplanted animals described in Example 6 is used for ELISA using an antibody specific for the synthetic FLAG epitope (IBI, Eastman Kodak, New Haven, CT) and compared against known standards of purified protein. Results indicate the presence of the anti-angiogenic polypeptide in the serum.

To determine whether a functional anti-angiogenesis polypeptide is present in the circulation, sera from transplanted animals is tested for its ability to inhibit the proliferation of bovine capillary cells *in vitro*; O'Reilly (1994). Briefly, cells are plated in 24 well dishes at 25,000 cells/ml and maintained in DMEM with 5% bovine calf serum for 24 hours. The medium is then replaced with fresh medium containing various dilutions of the test serum. After 20 minutes of incubation, fresh medium including b-FGF (final concentration 1 ng/ml) is added and the cells are cultured for 72 hours. Cells are then dispersed using trypsin and the cell number determined by Coulter counter. Results indicate that functional anti-angiogenic polypeptide is present in the sera of the recipient mice.

In addition, the ability of circulating anti-angiogenic polypeptide to inhibit the growth of primary tumor cells is assessed. Transplanted mice are subcutaneously injected with one million Lewis lung carcinoma (LLC) cells (O'Reilly, (1994)) at the proximal midline of their dorsal skin. The mice are closely monitored for survival, tumor size and growth, and overall health. Results indicate that the anti-angiogenic polypeptides from the sera of the

10

15

-26-

recipient mice inhibit growth of the LLC tumor cells.

Finally, upon sacrifice of the transplanted recipient mice, blood, spleen, thymus and bone marrow are harvested and analyzed for the presence of proviral DNA by Southern analysis as well as expression of the transferred GFP and anti-angiogenic polypeptide cDNAs by flow cytometry and ELISA. Moreover, a portion of bone marrow cells is re-transplanted into secondary recipients to generate individual day 12 spleen colonies, as well as plated in methylcellulose to assess in vitro clonogenic progenitors. Individual clones are analyzed for proviral DNA by PCR or Southern blot, and for gene expression by flow cytometry and ELISA. Results of these tests also indicate the presence of proviral DNA and expression of the anti-angiogenic polypeptides and marker proteins.

Example 10: Evaluating the Efficacy of Retroviral Gene Therapy Vectors

Encoding Anti-Angiogenic Polypetides on Various Human

Cancers Implanted in SCID Mice Using Ex Vivo Gene Therapy

This example illustrates a method for rapidly screening various forms of human cancer to determine susceptibility to treatment by the systemic delivery of anti-angiogenic polypeptides.

The methods for gene transfer, assessment of proviral marking and
assessment of transferred gene expression as described in Examples 3 through
9 are repeated using immuno-deficient SCID mice, with the following
exceptions. Since SCID mice are more sensitive to γ-irradiation than
C57BL6/J mice, the female SCID recipients receive a lower dose of 400cGy of
whole body irradiation in contrast to the 950cGy required for C57BL6/J. In
addition, since the SCID mice do not possess allelic differences at the CD45
cell surface antigen locus, donor and recipient cells are phenotypically
distinguished on the basis of Y chromosome specific sequences using Southern

blot analysis.

5

15

20

25...

Bone marrow from male donor SCID mice is infected, selected for on the basis of expression of the transferred GFP marker cDNA, and transplanted into irradiated female SCID recipients. Engraftment with provirally marked cells and expression of the transferred genes is demonstrated. The mice are then separately implanted with a variety of human tumor cell types, e.g., breast adenocarcinoma, lung squamous cell carcinoma, and brain glioblastoma. In each case, the ability of the anti-angiogenic polypeptides to inhibit the growth of the various human tumor cell types is monitored and quantified.

10 Example 11: Evaluating the Efficacy of Retroviral Gene Therapy Vectors

Encoding Anti-Angiogenic Polypeptides for Treatment of

Ovarian Cancer Using In Vivo Gene Therapy

This example illustrates the feasibility of using retroviral gene therapy vectors encoding anti-angiogenic polypeptides to achieve efficient gene transfer to established tumors in vivo using a well-established murine model of human ovarian cancer. Following injections, mice are closely monitored for tumor growth and survival.

Eight to ten week old nude mice (Jackson Labs, Bar Harbor are injected intra-peritoneally with 1 x 10⁷ PA-1 cells, an ovarian cancer cell-line (ATCC), and followed until palpable tumors are identified. Viral supernatant for <u>in vivo</u> injection is prepared as follows: Viral producer cells are grown to confluence in DMEM with 10% bovine calf serum, and the medium is then changed. After 24 hours of incubation, the viral conditioned supernatant is filtered though a 0.45 um low protein binding filter, protamine sulfate is added to a final concentration of 6ug/ml, the solution is aliquoted into 2 ml volumes, and frozen at -80°C. Recipient mice receive three intraperitoneal injections of viral supernatant (2 mls per injection) in addition to the polycation, over a period of

10

36 hours. Control mice are injected with medium collected from confluent dishes of NIH3T3 cells. Following injection of the viral conditioned supernatant, the mice are analyzed for survival as well as tumor growth over time as compared to mock injected controls. Results indicate that treatment of the ovarian cancer occurs. At death, the tumors are removed, weighed, and the cells dissociated for DNA extraction for Southern blot analysis to detect recombinant provirus.

Those skilled in the art will be able to ascertain using no more than routine experimentation, many equivalents of the specific embodiments of the invention described herein. These and all other equivalents are intended to be encompassed by the following claims.

In other embodiments, the invention provides methods and compositions for treating diseases and processes that are mediated by angiogenesis including, but not limited to, hemangioma, solid tumors, leukemia, metastasis,

15 telangiectasia, psoriasis, scleroderma, pyogenic granuloma, myocardial angiogenesis, plaque neovascularization, coronary collaterals, cerebral collaterals, arteriovenous malformations, ischemic limb angiogenesis, corneal diseases, rubeosis, neovascular glaucoma, diabetic retinopathy, retrolental fibroplasia, arthritis, diabetic neovascularization, macular degeneration, wound

20 healing, peptic ulcer, *Helicobacter* related diseases, fractures, keloids, vasculogenesis, hematopoiesis, ovulation, menstruation, placentation, and cat scratch fever.

What is claimed is:

10

15

20

-29-

CLAIMS

- 1. Use of a nucleic acid molecule which expresses an anti-angiogenic polypeptide selected from the group consisting of human angiostatin, murine angiostatin, human endostatin, murine endostatin, and angiogenesis-inhibiting fragments thereof in the preparation of a medicament for inhibiting tumor growth in a human patient harboring a solid tumor, wherein expression of the anti-angiogenic polypeptide in the patient inhibits angiogenesis in the vicinity of the tumor and/or systemically by diffusion of the recombinant protein to the vascular compartment from secreting transduced cells, thereby inhibiting its growth.
- 2. Use of a nucleic acid molecule which expresses an anti-angiogenic polypeptide of the formula A-B, wherein

A and B are polypeptide and/or export signal joined by a polypeptide and/or export signal bond;

A contains at least amino acids and comprises kringles 1, 2, and 3 of human or murine angiostatin; and

B contains at least amino acids and includes at least 75% of the amino acid sequence of human or murine endostatin in the preparation of a medicament for inhibiting tumor growth in a human patient harboring a solid tumor, wherein expression of the anti-angiogenic polypeptide in the patient inhibits angiogenesis in the vicinity of the tumor and/or systemically by diffusion of the recombinant protein to the vascular compartment from secreting transduced cells, thereby inhibiting its growth.

3. The use of claim 2, wherein A further comprises kringle region 4 of

human or murine angiostatin.

- 4. The use of claim 2 or claim 3, wherein A further comprises kringle 5 of human or murine plasminogen.
- 5. The use of claim 1 or claim 2, wherein said nucleic acid moleculeconstitutes a portion of a viral vector.
 - 6. The use of claim 1 or claim 2, wherein said nucleic acid molecule constitutes aportion of a plasmid.
 - 7. The use of claim 6, wherein said plasmid is carried in a cell-free carrier so that the plasmid transfects living cells of the patient following plasmid administration, causing expression of the anti-angiogenesis polypeptide and/or export signal in the patient such that angiogenesis in the vicinity of the tumor and/or systemically by diffusion of the recombinant protein to the vascular compartment from secreting transduced cells is inhibited, causing inhibition of tumor growth.
 - 8. The use of claim 6, wherein said plasmid has been transfected into animal cells *ex vivo*, wherein said animal cells express the anti-angiogenesis polypeptide to inhibit tumor-associated angiogenesis and tumor growth.
 - 9.* The use of claim 5, wherein said viral vector is carried in a cell-free carrier, so that the viral vector is incorporated into living cells of the patient following viral vector administration, causing expression of the anti-

20

10


angiogenesis polypeptide in the patient such that angiogenesis in the vicinity of the tumor and/or systemically by diffusion of the recombinant protein to the vascular compartment from secreting transduced cells is inhibited, causing inhibition of tumor growth.

- 10. The use of claim 5, wherein animal cells are infected with said viral vector *ex vivo* and then administered to said patient, wherein said animal cells express the anti-angiogenesis polypeptide to inhibit tumor-associated angiogenesis and tumor growth.
 - 11. The use of claim 8, wherein said animal cells are human cells.
- 10 12. The use of claim 11, wherein said human cells are autologous.
 - 13. The use of claim 11, wherein said human cells are allogeneic.
 - 14. The use of claim 10, wherein said animal cells are human cells.
 - 15. The use of claim 14, wherein said human cells are autologous.
 - 16. The use of claim 14, wherein said human cells are allogeneic.
- 15 17. The use of claim 5, wherein said viral vector is a retroviral vector.
 - 18: The use of claim 5, wherein said viral vector is a non-retroviral vector selected from the group consisting of adenoviral, adeno-associated, herpes, Simliki Forest virus, and poxvirus vectors.

- 19. The use of claim 17, wherein said retroviral vector is Murine Stem Cell Virus or a lentivirus.
- 20. The use of claim 1, wherein said angiostatin comprises kringles 1, 2 and 3.
- 5 21. The use of claim 20, wherein said angiostatin further comprises kringle 4.
 - 22. The use of claim 1, wherein said anti-angiogenic polypeptide is a fusion of angiostatin or a biologically active fragment thereof and endostatin or a biologically active fragment thereof.
- 10 23. The use of claim 1, wherein said nucleic acid molecule includes a nucleotide sequence encoding a signal polypeptide and/or export signal for effecting secretion of said anti-angiogenesis polypeptide.
 - 24. The use of claim 23, wherein said signal polypeptide and/or export signal is plasminogen signal polypeptide and/or export signal.
- 15 25. The use of claim 1, wherein said nucleic acid molecule includes a nucleotide sequence encoding a preactivation polypeptide and/or export signal for effecting Golgi and/or ER export of the anti-angiogenic polypeptide..
 - 26. The use of claim 25, wherein said preactivation polypeptide and/or export signal is a preactivation polypeptide and/or export signal of a human protein of the blood coagulation cascade.

- 27. The use of claim 26, wherein said preactivation polypeptide and/or export signal is human plasminogen preactivation polypeptide and/or export signal.
- 28. The method of claim 25, wherein the preactivation encoding sequence is positioned between a signal-encoding sequence and the sequence encoding the anti-angiogenic polypeptide and/or export signal.
 - 29. The use of claim 1, wherein said nucleic acid molecule includes a nucleotide sequence encoding a tag for identification of said anti-angiogenic polypeptide.
- 30. The method of claim 27, wherein said tag is a Flag tag polypeptide and/or export signal.
 - 31. A viral gene therapy vector comprising a nucleic acid molecule which encodes an anti-angiogenic polypeptide selected from the group consisting of human angiostatin, murine angiostatin, human endostatin, murine endostatin, and angiogenesis-inhibiting fusions and fragments thereof, wherein said viral vector is sufficiently attenuated for use in human gene therapy.
 - 32. A human cell infected with the vector of claim 31.
- 33. Use of a nucleic acid molecule which expresses in said patient an anti-angiogenic polypeptide selected from the group consisting of human
 angiostatin, murine angiostatin, human endostatin, murine endostatin, and angiogenesis-inhibiting fusions and fragments thereof, in the preparation of a

medicament for treating a human patient suffering from diabetic retinopathy, wherein expression of the anti-angiogenic polypeptide in the patient inhibits angiogenesis in the vicinity of the retina.



2/8





Fig. 2



WO 99/26480 PCT/US98/24950 _

4/8





77:1

5/8

ACTTGGATTGACCGAGTGATGAGAAATAATTAA ThrTrplleGluGlyValMetArgAsnAsn

rend human plasminagen coding sequence.

The state of the s

Fig 6

GATERIACTUT A Lader Leuffe

Signal peptide

260 TECENTATION A CHILLETINES

600 CACCCCCCCC ProProPro

Helashtylisys

SGO GLTACACTAC TyaThaTha

400 TRECUCKAÇI FheProHisVal

Kringles | Kringles | -3

TO ANTIFICIAL STATE | 110

TO ANTIFIC STATE | 110

TO ANTIFIC

100 ACTUTATOR Valty: Leu

EGG CTCCAAGACC SerlysThr

TAGET TIAGET TO THE TOTAL TO THE TOTAL THE TOT Preactivation peptide (PAP) Page Mou Plasm cDNA

Page 2

GTGCCAGTACT CysGInSer

J JOO GCTGACAAGG GlyasplysGly

1400 TGCCCACAGT ProThrVal Profit poly of the control of the co

Trocheant $\ker_{\mathcal{A}} = \ker_{\mathcal{A}} =$

AGCTEGCACC AlaGlyThr

SIT

1600 TGCCGAAACC CysArgAsnPro

CATCCTITICA Sor PheGlu

1500 AGACCTGAAT Argi's oGluphe 1800 CAGCCTTAGA Sex Lewlurg

ئ

BNSDOCID: <WO__9926480A1_I_>

(Sheet 1 of

Tue, Oct 21, 1997 Code: Universal			1	Mou Plasm c	DNA			Page
2010	2020	2030	2040	2050	2060	2070	2080	2090
CATTGCCCTGCTGA IleAlaLeuLeuL	AACTAAGCC ysLeuSerA	GCCCAGCCAC rgProAlaTh	CATCACGGAT rIleThrAsp	AAAGTCATTC LysVallleP	CAGCTTGTC	CCCATCTCCAA euProSerProA	ATTACATGG snTyrMetV	MGCIGA alAlaAsp
2190 CCGGACAATA ArgThrile								
2110	2120	2130	2140	2150	2160	2170	2180	2190
TGTTACATCACCGG CysTyrlleThrGl	CTGGGGAGA YTTpGlyGl	GACTCAAGGG uThrGlnGly	ACTTICGGIG ThrPheGlyA	CCGGTCGTCT laGlyArgLe	CAAGGAGGC uLysGluAl	CAGCIGCCIGI aGlnLeuProVa	lileGluAs	nLysVal
2200 TGCAACCGCG CysAsnArgVal								
2210 TCGAGTATCTGAAC GluTyrLeuAsn	2220 AACAGAGTC AsnArgVal	2230 AAATCCACGG LysSerThrG	2240 AGCTCTGTGC luLeuCysAl	2250 CGGGCAACTG aGlyGlnLeu	2260 GCTGGTGGC AlaGlyGly	2270 STCGACAGCTGC ValAspSerCys	2280 CAAGGCGAC GlnGlyAsp	2290 AGTGGAG SerGlyGly
2300 GACCTCTGGT ProLeuVal								
2310	2320	2330	2340	2350	2360	2370	2380	2390
TTGCTTCGAGAAGG CyaPheGluLysA	ACAAGTACA spLysTyrI	TTTTACAAGG leLeuGlnGl	yValThrSer	TrpGlyLeuG	lyCysAlaA	rgProAsnLysP	rociyvain	yrvalArg
2400 TGTCTCACGG			end	plasmi	nogen	coding	seque	ence.
ValSerArg								
2410 TTTGTTGATTGGAT PheValAspTrpIl	2420 TGAAAGGGA eGluArgGl	2430 GATGAGGAAT uMetArgAsn	2140 AACTGACTAC ASTERCLEUC	2450 GTGGAAGGCC :lyGlyArgPr	2460 GAGCAAAAC OSerLysTh	2470 CTCTGCTTACTA rSerAlaTyrEn	2480 AAGCTTACT dSerLeuLe	2490 GAATATG uAsnMet
2500 GGGAGAGGGC GlyArgGlyLeu								
2510 TTAGGGTGTTTGGA ArgValPheGly	2520 AAAACTGAC LysThrAsp	2530 AGTAATCAAA SerAsnGlnT	2540 CTGGGACACT hrGlyThrLe	2550 ACACTGAACC wHisEndThr	2560 ACAGCTTCC ThrAlaSer	2570 TGTCGCCCCTCA CysArgProSer	2580 GCCCCTCCC AlaProPro	2590 CTTTTT LeuPheLeu
2600 TGTATTATTG TYTTYTCYS								
2610 TGGGTAAAATTTTC GlyEndAsnPheF	2620 CTGTCTGTG ProValCysG	2630 GACTTCTGGA lyLeuLeuAs	2640 TTTTGTGACA pPheValThr	2650 ATAGACCATO IleAspHish	2660 ACTGCTGTG LisCysCysA	2670 ACCTTTGTTGAA spLeuCysEndL	2680 AATAAACTC ysEndThrA	2690 GATACTT rgTyrLeu
2700 ACTTIG Leu								

SEQUENCE LISTING

<110> Genetix Pharmaceuticals, Inc.

<120> ANTI-ANGIOGENIC GENE THERAPY VECTORS AND
THEIR USE IN TREATING ANGIOGENISIS-RELATED DISEASES

<130>	50033/002WO1
-1140>	PCT/US98/xxxxx
	1998-11-20
(111)	1330 11 20
<150>	08/975,424
<151>	1997-11-20
<160>	20
<170>	FastSEQ for Windows Version 3.0
<210>	
<211>	
<212>	
<213>	Homo sapiens
<220>	
<221>	
	(1)(2433)
12227	(2, (2.25)
<400>	1
atg gaa cat	aag gaa gtg gtt ctt cta ctt ctt tta ttt ctg aaa tca 48
Met Glu His	Lys Glu Val Val Leu Leu Leu Leu Phe Leu Lys Ser
1	5 10 15
aat aaa aaa	gag cct ctg gat gac tat gtg aat acc cag ggg gct tca 96
	Glu Pro Leu Asp Asp Tyr Val Asn Thr Gln Gly Ala Ser
Gry Grn Gry	20 25 30
	20 23 30
ctg ttc agt	gtc act aag aag cag ctg gga gca gga agt ata gaa gaa 144
Leu Phe Ser	Val Thr Lys Lys Gln Leu Gly Ala Gly Ser Ile Glu Glu
35	40 45
	aaa tgt gag gag gac gaa gaa ttc acc tgc agg gca ttc 192
_	Lys Cys Glu Glu Asp Glu Glu Phe Thr Cys Arg Ala Phe
50	55 60
caa tat cac	agt aaa gag caa caa tgt gtg ata atg gct gaa aac agg 240
	Ser Lys Glu Gln Gln Cys Val Ile Met Ala Glu Asn Arg
65	70 75 80
	4
aag tcc tcc	ata atc att agg atg aga gat gta gtt tta ttt gaa aag 288
Lys Ser Ser	Ile Ile Ile Arg Met Arg Asp Val Val Leu Phe Glu Lys
	85 90 95

	gtg Val															336
	acg Thr															384
	act Thr 130				_		_				_					432
	gga Gly	-														480
	ccc Pro															528
	att Ile															576
	gac Asp					_		_			_	_	-	_	_	624
	gac Asp 210															672
	aac Asn	_		_	_	_			_	_			_			720
_	cgg Arg			_				_			_	_				768
_	gac. Asp			_	_											816
	cag Gln	_	_	_				_			_				_	864
_	acc Thr 290	Val		_					His			Ala				912
	aca Thr															960

														tgg Trp			1008
į	aca Thr	acc Thr	aac Asn	agc Ser 340	caa Gln	gtg Val	cgg Arg	tgg Trp	gag Glu 345	tac Tyr	tgt Cys	aag Lys	ata Ile	ccg Pro 350	tcc Ser	tgt Cys	1056
														gca Ala			1104
														gga Gly			1152
•														tgt Cys			1200
														gaa Glu			1248
														gat Asp 430			1296
														tgg Trp			1344
														gta Val			1392
		Pro	Val	Val	Leu							Pro		gaa Glu			1440
														gcg Ala			1488
	gtt Val	act Thr	ggg Gly	acg Thr 500	cca Pro	tgc Cys	cag Gln	gac Asp	tgg Trp 505	gct Ala	gcc Ala	cag Gln	gag Glu	ccc Pro 510	cat His	aga Arg	1536
														ctg Leu			1584 ************************************
														tgg Trp			1632

530		535	5	540	
_	-	_		gat gtc cct cag Asp Val Pro Gln	_
				gtg gag ccg aag Val Glu Pro Lys 575	
_			ys Val Ala H	eac cca cat tcc His Pro His Ser 590	
	_	-		tg cac ttc tgt let His Phe Cys 605	
			al Leu Thr A	ct gcc cac tgc la Ala His Cys 20	=
				tc ctg ggt gca le Leu Gly Ala	
				ta gaa gtg tct le Glu Val Ser 655	
-			sp Ile Ala L	tg cta aag cta eu Leu Lys Leu 670	
	_	-	_	ct tgt ctg cca la Cys Leu Pro 685	
			nr Glu Cys P	tc atc act ggc he Ile Thr Gly 00	
				tc aag gaa gcc eu Lys Glu Ala	
				at gag ttt ctg yr Glu Phe Leu 735	
			s Ala Gly H	at ttg gcc gga is Leu Ala Gly 750	
act gac agt	tgc cag ggt	gac agt gg	ga ggt cct c	tg gtt tgc ttc	gag 2304

	Thr	Asp	Ser 755	Cys	Gln	Gly	Asp	Ser 760	Gly	Gly	Pro	Leu	Val 765	Cys	Phe	Glu	
•	aag Lys	gac Asp 770	aaa Lys	tac Tyr	att Ile	tta Leu	caa Gln 775	gga Gly	gtc Val	act Thr	tct Ser	tgg Trp 780	ggt Gly	ctt Leu	ggc Gly	tgt Cys	2352
										gtt Val							2400
										aat Asn 810							2433

<210> 2 <211> 810 <212> PRT

<213> Homo sapiens

<400> 2 Met Glu His Lys Glu Val Val Leu Leu Leu Leu Phe Leu Lys Ser 10 Gly Gln Gly Glu Pro Leu Asp Asp Tyr Val Asn Thr Gln Gly Ala Ser 25 Leu Phe Ser Val Thr Lys Lys Gln Leu Gly Ala Gly Ser Ile Glu Glu 40 Cys Ala Ala Lys Cys Glu Glu Asp Glu Glu Phe Thr Cys Arg Ala Phe Gln Tyr His Ser Lys Glu Gln Gln Cys Val Ile Met Ala Glu Asn Arg Lys Ser Ser Ile Ile Ile Arg Met Arg Asp Val Val Leu Phe Glu Lys 90 Lys Val Tyr Leu Ser Glu Cys Lys Thr Gly Asn Gly Lys Asn Tyr Arg 105 100 Gly Thr Met Ser Lys Thr Lys Asn Gly Ile Thr Cys Gln Lys Trp Ser 120 115 Ser Thr Ser Pro His Arg Pro Arg Phe Ser Pro Ala Thr His Pro Ser 140 135 Glu Gly Leu Glu Glu Asn Tyr Cys Arg Asn Pro Asp Asn Asp Pro Gln 150 Gly Pro Trp Cys Tyr Thr Thr Asp Pro Glu Lys Arg Tyr Asp Tyr Cys 170 Asp Ile Leu Glu Cys Glu Glu Glu Cys Met His Cys Ser Gly Glu Asn 185 180 Tyr Asp Gly Lys Ile Ser Lys Thr Met Ser Gly Leu Glu Cys Gln Ala 200 205 Trp Asp Ser Gln Ser Pro His Ala His Gly Tyr Ile Pro Ser Lys Phe 220 215 ProgAsn: Lys. Asn Leu Lys. Lys. Asn. Tyr. Cys Arg Asn Pro Asp Arg Glu 230 235 Leu Arg Pro Trp Cys Phe Thr Thr Asp Pro Asn Lys Arg Trp Glu Leu 250 Cys Asp Ile Pro Arg Cys Thr Thr Pro Pro Pro Ser Ser Gly Pro Thr

			260					265					270		
Tyr	Gln	Cys 275		Lys	Gly	Thr	Gly 280			Tyr	Arg	Gly 285			Ala
Val	Thr 290	Val	Ser	Gly	His	Thr 295	Cys	Gln	His	Trp	Ser 300	Ala	Gln	Thr	Pro
His 305	Thr	His	Asn	Arg	Thr 310		Glu	Asn	Phe	Pro 315	Cys	Lys	Asn	Leu	Asp 320
Glu	Asn	Tyr	Cys	Arg 325	Asn	Pro	Asp	Gly	Lys 330	Arg	Ala	Pro	Trp	Cys 335	His
	Thr		340					345	_	_	_		350		_
_	Ser	355					360					365			
	Leu 370					375			_		380		_		
385	Arg	_			390					395	_	_	_		400
_	Ser			405					410	_				415	-
	Asn		420					425					430		_
_	Gly	435					440					445			_
_	Asn 450		_		-	455	_				460				
465	Pro				470		_			475					480
	Met		_	485			_		490	_				495	
	Thr	_	500					505					510		
	Ser Tyr	515					520			_		525			_
	530 Thr		_			535				_	540				_
545		11011	110		550		- 7 -	TIDP	- 7 -	555	· · · · ·			0111	560
	Ala			565	_	_	_	-	570					575	-
	Pro	_	580					585					590		
	Trp	595					600					605			
_	Thr 610					615	-				620				
625	Lys			_	630			_	-	635			_		640
	Glu			645					650					655	
-	Phe		660				-	665	F 4		•		6.70		. 6≥. €
	Pro	675					680					685			
Pro	Asn 690	TYT	val	vai	Ala	Asp 695	Arg	Thr	GIu	Cys	700	тте	Thr	GIY	ırp

										-						
Gly 705	Glu	Thr	Gln	Gly	Thr 710	Phe	Gly	Ala	Gly	Leu 715	Leu	Lys	Glu	Ala	Gln 720	
			Ile	725					730					735		
-	_		Gln 740					745					750			
	_	755	Cys				760					765				
_	770		Tyr			775					780					
785			Asn		790					Arg 795	Val	Ser	Arg	Phe	Val 800	
Thr	Trp	Ile	Glu	Gly 805	Val	Met	Arg	Asn	Asn 810							
	<2 <2	212>	2696		culus	3										
		220>	GD C													
			(1)	(2	2436)	1										
		00>									4 - 4					4.0
			aag Lys													48
gga Gly	caa Gln	gly aaa	gac Asp 20	tcg Ser	ctg Leu	gat Asp	ggc Gly	tac Tyr 25	ata Ile	agc Ser	aca Thr	caa Gln	30 Gly 39	gct Ala	tca Ser	96
			ctc Leu													144
tgt Cys	ttg Leu 50	gcc Ala	aaa Lys	tgt Cys	gaa Glu	999 Gly 55	gaa Glu	aca Thr	gac Asp	ttt Phe	gtc Val 60	tgc Cys	agg Arg	tca Ser	ttc Phe	192
cag Gln 65	tac Tyr	cac His	agc Ser	aaa Lys	gag Glu 70	cag Gln	caa Gln	tgc Cys	gtg Val	atc Ile 75	atg Met	gcg Ala	gag Glu	aac Asn	agc Ser 80	240
			tcc Ser													288
			ctg Leu 100												aga Argasasse	336
gga Gly	acc Thr	atg Met	tcc Ser	agg Arg	aca Thr	aag Lys	agt Ser	ggt Gly	gtt Val	gcc Ala	tgt Cys	caa Gln	aag Lys	tgg Trp	ggt Gly	384

Line to the state of

																_	_
		115					120					125	į				-
gcc Ala	acg Thr	Phe	ccc Pro	Cac	gta Val	ccc Pro 135	aac Asn	tac Tyr	tct Ser	ccc	agt Ser 140	Thr	cat His	ccc Pro	aat Asn	432	
gag Glu 145	Gly	cta Leu	gaa Glu	gag Glu	aac Asn 150	tac Tyr	tgt Cys	agg Arg	aac Asn	cca Pro 155	Asp	aat Asn	gat Asp	gaa Glu	caa Gln 160	480	
Gly aaa	cct Pro	tgg Trp	tgc Cys	tac Tyr 165	act Thr	aca Thr	gat Asp	ccg Pro	gac Asp 170	aag Lys	aga Arg	tat Tyr	gac Asp	tac Tyr 175	tgc Cys	528	
aac Asn	att Ile	cct Pro	gaa Glu 180	tgt Cys	gaa Glu	gag Glu	gaa Glu	tgc Cys 185	atg Met	tac Tyr	tgc Cys	agt Ser	gga Gly 190	Glu	aag Lys	576	
tat Tyr	gag Glu	ggc Gly 195	aaa Lys	atc Ile	tcc Ser	aag Lys	acc Thr 200	atg Met	tct Ser	gga Gly	ctt Leu	gac Asp 205	tgc Cys	cag Gln	gcc Ala	624	
tgg Trp	gat Asp 210	tct Ser	cag Gln	agc Ser	cca Pro	cat His 215	gct Ala	cat His	gga Gly	tac Tyr	atc Ile 220	cct Pro	gcc Ala	aaa Lys	ttt Phe	672	
cca Pro 225	agc Ser	aag Lys	aac Asn	ctg Leu	aag Lys 230	atg Met	aat Asn	tat Tyr	tgc Cys	cac His 235	aac Asn	cct Pro	gac Asp	gly aaa	gag Glu 240	720	
cca Pro	agg Arg	ccc Pro	tgg Trp	tgc Cys 245	ttc Phe	aca Thr	aca Thr	gac Asp	ccc Pro 250	acc Thr	aaa Lys	cgc Arg	tgg Trp	gaa Glu 255	tac Tyr	768	
tgt Cys	gac Asp	atc Ile	ccc Pro 260	cgc Arg	tgc Cys	aca Thr	aca Thr	ccc Pro 265	ccg Pro	ccc Pro	cca Pro	ccc Pro	agc Ser 270	cca Pro	acc Thr	816	
tac Tyr	caa Gln	tgt Cys 275	ctg Leu	aaa Lys	gga Gly	aga Arg	ggt Gly 280	gaa Glu	aat Asn	tac Tyr	cga Arg	999 Gly 285	acc Thr	gtg Val	tct Ser	864	
gtc Val	acc Thr 290	gtg Val	tct Ser	Gly 999	aaa Lys	acc Thr 295	tgt Cys	cag Gln	cgc Arg	tgg Trp	agt Ser 300	gag Glu	caa Gln	acc Thr	cct Pro	912	
cat His 305	agg Arg	cac His	aac Asn	agg Arg	aca Thr 310	cca Pro	gaa Glu	aat Asn	ttc Phe	ccc Pro 315	tgc Cys	aaa Lys	aat Asn	ctg Leu	gaa Glu 320	960	
gag Glu	aac Asn	tac Tyr	tgc, Cys	.¢gg. Arg 325	aac Asn	cca Pro	gat Asp	gga Gly	gaa Glu 330	act. Thr	gct. Ala	ccc Pro	tgg Trp	t gs Cys 335	t at . e Tyr	1008	
acc	act	gac	agc	cag	ctg	agg	tgg	gag	tac	tgt	gag	att	cca	tcc	tgc	1056	

	Thr	Thr	Asp	Ser 340	Gln	Leu	Arg	Trp	Glu 345	Tyr	Cys	Glu	Ile	Pro 350	Ser	Cys		
				_			gac Asp										13	104
							cag Gln 375										1,3	152
							act Thr										12	200
							cac His										12	248
							atg Met										12	296
	_				_		acc Thr		_	_	_	_					13	344
	_		_				tca Ser 455										13	92
							cca Pro										14	14 Ò
							aaa Lys										14	188
							cag Gln										15	36
							cag Gln										15	84
							gat Asp 535										16	32
•	aca				-		ctt Leu										16	80

_		_	tca Ser				-				_		_		_	1728
			999 Gly 580													1776
			caa Gln		_		-		-				_			1824
_			act Thr			_				_	_		-	_		1872
-	_		aaa Lys			_		_			_	-				1920
		_	gaa Glu			_		_	-	_	_	_			_	1968
_		_	atc Ile 660	_					_	_			_	_		2016
			cca Pro													2064
			aat Asn		_	_	_	_				_				2112
			gag Glu									_				2160
			Pro										_	_		2208
_			aga Arg 740	_			_			_	_					2256
		_	gac Asp	-	_			_	-				_	_	_	2304
		-	gac: Asp	_						_					ctt Leu	-√ - √2352

ggc Gly 785	tgt Cys	gct Ala	cgc Arg	ccc Pro	aat Asn 790	aag Lys	cct Pro	ggt Gly	gtc Val	tac Tyr 795	gtt Val	cgt Arg	gtc Val	tca Ser	cgg Arg 800	2400	
	gtt Val											tga	ctag	gtg		2446	
tgt gcc ttc	ttgg:	aaa a agc o ttt t	acto	gacag	gt aa ct ti	atcaa ctttg	aacto gtatt	g gga	acact tgtg	taca ggta	ctga aaat	aacca tttt	aca g	gctto	ttaggg cetgte gtggae ctegat	2506 2566 2626 2686 2696	
	< 2 < 2	210> 211> 212> 213>	812 PRT	muso	culus	5											
Met	Asp	400> His			Val	Ile	Leu	Leu		Leu	Leu	Leu	Leu		Pro		
1 Gly	Gln	Gly		5 Ser	Leu	Asp	Gly		10 Ile	Ser	Thr	Gln	Gly 30	15 Ala	Ser		
Leu	Phe	Ser 35	20 Leu	Thr	Lys	Lys	Gln 40	25 Leu	Ala	Ala	Gly	Gly 45		Ser	Asp		
Cys	Leu 50		Lys	Cys	Glu	Gly 55		Thr	Asp	Phe	Val 60		Arg	Ser	Phe		
Gln 65	Tyr	His	Ser	Lys	Glu 70		Gln	Cys	Val	Ile 75	Met	Ala	Glu	Asn	Ser 80		
Lys	Thr			85					90					95			
_	Val	_	100					105					110				
_	Thr	115					120					125					
	Thr 130					135					140						
Glu 145	Gly	Leu	GIu	GIU	150	Tyr	Cys	Arg	Asn	155	Asp	ASI	Asp	GIU	160		
_	, P.ro	_		165					170					175	•		
	Ile		180					185					190				
_	Glu	195					200					205					
	Asp 210					215					220						
	Ser	Lys	Asn	Leu		Met	Asn	Tyr	Cys		Asn	Pro	Asp	Gly			
225 Dro	Arg	Pro	Trr	Cvs	230 Phe	Thr	Thr	Asn	Pro	235 Thr	Lvs	Ara	Trp	Glu	240 Tvr		
**	7.7			245	1110				250		-,-	3		355		4.	
Cys	Asp	Ile	Pro	Arg	Cys	Thr			Pro			Pro	Ser	Pro	Thr		

265

260

Tyr Gln Cys Leu Lys Gly Arg Gly Glu Asn Tyr Arg Gly Thr Val Ser 275 280 Val Thr Val Ser Gly Lys Thr Cys Gln Arg Trp Ser Glu Gln Thr Pro 295 300 His Arg His Asn Arg Thr Pro Glu Asn Phe Pro Cys Lys Asn Leu Glu 310 315 Glu Asn Tyr Cys Arg Asn Pro Asp Gly Glu Thr Ala Pro Trp Cys Tyr 325 330 Thr Thr Asp Ser Gln Leu Arg Trp Glu Tyr Cys Glu Ile Pro Ser Cys 345 Glu Ser Ser Ala Ser Pro Asp Gln Ser Asp Ser Ser Val Pro Pro Glu 360 Glu Gln Thr Pro Val Val Gln Glu Cys Tyr Gln Ser Asp Gly Gln Ser 375 Tyr Arg Gly Thr Ser Ser Thr Thr Ile Thr Gly Lys Lys Cys Gln Ser 390 395 Trp Ala Ala Met Phe Pro His Arg His Ser Lys Thr Pro Glu Asn Phe 410 Pro Asp Ala Gly Leu Glu Met Asn Tyr Cys Arg Asn Pro Asp Gly Asp 420 425 Lys Gly Pro Trp Cys Tyr Thr Thr Asp Pro Ser Val Arg Trp Glu Tyr 440 Cys Asn Leu Lys Arg Cys Ser Glu Thr Gly Gly Ser Val Val Glu Leu 455 Pro Thr Val Ser Gln Glu Pro Ser Gly Pro Ser Asp Ser Glu Thr Asp 470 475 Cys Met Tyr Gly Asn Gly Lys Asp Tyr Arg Gly Lys Thr Ala Val Thr 485 490 Ala Ala Gly Thr Pro Cys Gln Gly Trp Ala Ala Gln Glu Pro His Arg 500 505 His Ser Ile Phe Thr Pro Gln Thr Asn Pro Arg Ala Asp Leu Glu Lys 520 Asn Tyr Cvs Arg Asn Pro Asp Gly Asp Val Asn Gly Pro Trp Cys Tyr 535 540 Thr Thr Asn Pro Arg Lys Leu Tyr Asp Tyr Cys Asp Ile Pro Leu Cys 550 555 Ala Ser Ala Ser Ser Phe Glu Cys Gly Lys Pro Gln Val Glu Pro Lys 570 Lys Cys Pro Gly Arg Val Val Gly Gly Cys Val Ala Asn Pro His Ser 585 Trp Pro Trp Gln Ile Ser Leu Arg Thr Arg Phe Thr Gly Gln His Phe 595 600 Cys Gly Gly Thr Leu Ile Ala Pro Glu Trp Val Leu Thr Ala Ala His 615 620 Cys Leu Glu Lys Ser Ser Arg Pro Glu Phe Tyr Lys Val Ile Leu Gly 630 635 Ala His Glu Glu Tyr Ile Arg Gly Leu Asp Val Gln Glu Ile Ser Val 650 645 Ala Lys Leu Ile Leu Glu Pro Asn Asn Arg Asp Ile Ala Leu Leu Lys 665 Leu Ser Arg Pro Ala Thr Ile Thr Asp Lys Val Ile Pro Ala Cys Leu 680 Fre Ser Prov**Ast Tyr. Met** Wal Ala Asp Args That Ilea Cys Tyr Ile Thr 🐝 🕬 🔻 Gly Trp Gly Glu Thr Gln Gly Thr Phe Gly Ala Gly Arg Leu Lys Glu

705					710					715					720		
Ala	Gln	Leu	Pro	Val 725	Ile	Glu	Asn	Lys	Val 730	Cys	Asn	Arg	Val	Glu 735	Tyr		
Leu	Asn	Asn	Arg 740	Val	Lys	Ser	Thr	Glu 745	Leu	Cys	Ala	Gly	Gln 750	Leu	Ala		
Gly	Gly	Val 755		Ser	Cys	Gln	Gly 760		Ser	Gly	Gly	Pro 765	Leu	Val	Cys		
Phe	Glu		Asp	Lys	Tyr			Gln	Gly	Val	Thr 780		Trp	Gly	Leu		
•	770 Cys	Ala	Arg	Pro		775 Lys	Pro	Gly	Val			Arg	Val	Ser			
785 Phe	Val	Asp	Trp		790 Glu	Arg	Glu	Met		795 Asn	Asn				800		
				805					810								
		210>															
			1083	3													
		212>			ni en	-											
	< 4	213>	ноше	sar	Tell	•											
	<2	220>															
	<2	221>	CDS														
	<:	222>	(1)	(]	1083)											
		400>															
	ctc															4	18
-	Leu	Ser	Glu		Lys	Thr	Gly	Asn		Lys	Asn	Tyr	Arg		Thr		
1				5					10					15			
atq	tcc	aaa	aca	aaa	aat	aac	atc	acc	tat	caa	aaa	tgg	agt	tcc	act	ç	96
	Ser																
			20					25					30				
									~~+	202	a.a.	~~~	t 02	asa	aa a	7.4	14
	ccc Pro															4.7	**
ser	PIO	35	Arg	FLO	AL G	FIIC	40	110	nia	1111	****	45	001		1		
		•															
	gag															19	92
Leu	Glu	Glu	Asn	Tyr	Cys		Asn	Pro	Asp	Asn		Pro	Gln	Gly	Pro		
	50					55					60						
taa	tgc	tat	act	act	gat.	сса	gaa	aaq	aσa	tat	gac	tac	tac	gac	att	24	40
	Cys																
65			-		70			-	_	75	_	_			80		
																2.0	
	gag Glu															20	38
rea	GIU	Cys	GIU	85	GIU	Cys	Mec	птэ	90	261	GIY	GIU	ASII	95	nop		
ggc	aaa	att	tcc	aag	acc	atg	tct	gga	ctg	gaa	tgc	cag	gcc	tgg	gac	33	36
Gly	Lys	Ile		Lys	Thr	Met	Ser		Leu	Glu	Cys	Gln		Trp	Asp		
			100					105					110				
+~+	cag	age	רר פּ		get	. cat	- നമ ല	tac	ant	ocr.	√6 000	ക്രാകം	ttt.	. cca	aac	31	8 4. 256 (1990) 1990)
Ser	Gln	Ser	Pro	His	Ala	His	Glv	Tvr	Ile	Pro	Ser	Lys	Phe	Pro	Asn		THE STATE OF THE S
							120					125					

120

115

_		ctg Leu	_	_			_	_			_					432
		tgt Cys				_			_	_		_			-	480
		cgc Arg														528
		aag Lys														576
		999 Gly 195														624
		agg Arg			_				-			_	_	_		672
	_	cgc Arg			_				-		-					720
	-	caa Gln														768
		gta Val		_	-		_	_			_					816
		gtg Val 275														864
		tcc Ser							_	_	_	-				912
	_	aca Thr					-	_			_					960
_		ctg Leu		_			_				-	-				1008
		tgt Cys														1056

240 345 350

Ctg aaa aaa tgc tca gga aca gaa gcg 1083

Leu Lys Lys Cys Ser Gly Thr Glu Ala

<210> 6 <211> 361 <212> PRT <213> Homo sapiens

(213) 110110 5451011

355

<400> 6 Tyr Leu Ser Glu Cys Lys Thr Gly Asn Gly Lys Asn Tyr Arg Gly Thr 10 Met Ser Lys Thr Lys Asn Gly Ile Thr Cys Gln Lys Trp Ser Ser Thr 20 25 Ser Pro His Arg Pro Arg Phe Ser Pro Ala Thr His Pro Ser Glu Gly 40 Leu Glu Glu Asn Tyr Cys Arg Asn Pro Asp Asn Asp Pro Gln Gly Pro 55 Trp Cys Tyr Thr Thr Asp Pro Glu Lys Arg Tyr Asp Tyr Cys Asp Ile 70 75 Leu Glu Cys Glu Glu Glu Cys Met His Cys Ser Gly Glu Asn Tyr Asp 85 90 Gly Lys Ile Ser Lys Thr Met Ser Gly Leu Glu Cys Gln Ala Trp Asp 100 105 Ser Gln Ser Pro His Ala His Gly Tyr Ile Pro Ser Lys Phe Pro Asn 115 120 Lys Asn Leu Lys Lys Asn Tyr Cys Arg Asn Pro Asp Arg Glu Leu Arg 135 Pro Trp Cys Phe Thr Thr Asp Pro Asn Lys Arg Trp Glu Leu Cys Asp 150 155 Ile Pro Arg Cys Thr Thr Pro Pro Pro Ser Ser Gly Pro Thr Tyr Gln 170 165 Cys Leu Lys Gly Thr Gly Glu Asn Tyr Arg Gly Asn Val Ala Val Thr 185 190 180 Val Ser Gly His Thr Cys Gln His Trp Ser Ala Gln Thr Pro His Thr 200 His Asn Arg Thr Pro Glu Asn Phe Pro Cys Lys Asn Leu Asp Glu Asn 215 220 Tyr Cys, Arg. Asn. Pro. Asp. Gly, Lys Arg. Ala. Pro. Trp. Cys. His. Thr. Thr 230 235 Asn Ser Gln Val Arg Trp Glu Tyr Cys Lys Ile Pro Ser Cys Asp Ser 250 245 Ser Pro Val Ser Thr Glu Gln Leu Ala Pro Thr Ala Pro Pro Glu Leu 265 260 Thr Pro Val Val Gln Asp Cys Tyr His Gly Asp Gly Gln Ser Tyr Arg 280 Gly Thr Ser Ser Thr Thr Thr Gly Lys Lys Cys Gln Ser Trp Ser 300 295 Ser Met Thr Pro His Arg His Gln Lys Thr Pro Glu Asn Tyr Pro Asn 305 4 5 310 310 3335 4 Ala Gly Leu Thr Met Asn Tyr Cys Arg Asn Pro Asp Ala Asp Lys Gly 330 325

			Phe 340 Cys					Ser 345 Ala	Val	Arg	Trp	Glu	Tyr 350	Cys	Asn	
	<2 <2	210> 211> 212>	108		culus	5										
	<2	220> 221> 222>	CDS	(:	1086)										
	tat		tca					ggc Gly								48
	_							gtt Val 25								96
_				_				tct Ser		_						144
								aac Asn								192
		_				-	_	gac Asp	_	_		_				240
		_	_	-		_	_	atg Met		_	_					288
					_		_	tct Ser 105			-					336
_		_	_			-		gga Gly				_				384
_	_		_	_	_			tgc Cys				_				432
			_				_	ecc Pro			_		_		tgt Cys ^{c.} 160	480

	atc Ile															52	28
	tgt Cys															57	76
	gtg Val															62	24
	cac His 210															67	72
aac Asn 225	tac Tyr	tgc Cys	cgg Arg	aac Asn	cca Pro 230	gat Asp	gga Gly	gaa Glu	act Thr	gct Ala 235	ccc Pro	tgg Trp	tgc Cys	tat Tyr	acc Thr 240	72	20
act Thr	gac Asp	agc Ser	cag Gln	ctg Leu 245	agg Arg	tgg Trp	gag Glu	tac Tyr	tgt Cys 250	gag Glu	att Ile	cca Pro	tcc Ser	tgc Cys 255	gag Glu	76	58
tcc Ser	tca Ser	gca Ala	tca Ser 260	cca Pro	gac Asp	cag Gln	tca Ser	gat Asp 265	tcc Ser	tca Ser	gtt Val	cca Pro	cca Pro 270	gag Glu	gag Glu	83	16
caa Gln	aca Thr	cct Pro 275	gtg Val	gtc Val	cag Gln	gaa Glu	tgc Cys 280	tac Tyr	cag Gln	agc Ser	gat Asp	999 Gly 285	cag Gln	agc Ser	tat Tyr	86	64
	ggt Gly 290															91	12
	gct Ala															96	60
	gct Ala		Leu.			Asn	Tyr		Arg,							100	38
	cct Pro															109	56
	ctg Leu															108	86

<210> 8 <211> 362

<212> PRT

<212> DNA

<213> Mus musculus

```
<213> Mus musculus
     <400> 8
Val Tyr Leu Ser Glu Cys Lys Thr Gly Ile Gly Asn Gly Tyr Arg Gly
                              10
Thr Met Ser Arg Thr Lys Ser Gly Val Ala Cys Gln Lys Trp Gly Ala
                            25
    20
Thr Phe Pro His Val Pro Asn Tyr Ser Pro Ser Thr His Pro Asn Glu
35 40
Gly Leu Glu Glu Asn Tyr Cys Arg Asn Pro Asp Asn Asp Glu Gln Gly
                   55
Pro Trp Cys Tyr Thr Thr Asp Pro Asp Lys Arg Tyr Asp Tyr Cys Asn
              70
                                75
Ile Pro Glu Cys Glu Glu Glu Cys Met Tyr Cys Ser Gly Glu Lys Tyr
             85
                        90
Glu Gly Lys Ile Ser Lys Thr Met Ser Gly Leu Asp Cys Gln Ala Trp
                           105
Asp Ser Gln Ser Pro His Ala His Gly Tyr Ile Pro Ala Lys Phe Pro
                       120
 115
Ser Lys Asn Leu Lys Met Asn Tyr Cys His Asn Pro Asp Gly Glu Pro
                    135
                           140
Arg Pro Trp Cys Phe Thr Thr Asp Pro Thr Lys Arg Trp Glu Tyr Cys
                     155
                150
Asp Ile Pro Arg Cys Thr Thr Pro Pro Pro Pro Pro Ser Pro Thr Tyr
                  170 175
             165
Gln Cys Leu Lys Gly Arg Gly Glu Asn Tyr Arg Gly Thr Val Ser Val
                            185
Thr Val Ser Gly Lys Thr Cys Gln Arg Trp Ser Glu Gln Thr Pro His
                        200
Arg His Asn Arg Thr Pro Glu Asn Phe Pro Cys Lys Asn Leu Glu Glu
                   215
                                     220
Asn Tyr Cys Arg Asn Pro Asp Gly Glu Thr Ala Pro Trp Cys Tyr Thr
       230 . 235
Thr Asp Ser Gln Leu Arg Trp Glu Tyr Cys Glu Ile Pro Ser Cys Glu
             245
                              250
Ser Ser Ala Ser Pro Asp Gln Ser Asp Ser Ser Val Pro Pro Glu Glu
                            265
                                             270
Gln Thr Pro Val Val Gln Glu Cys Tyr Gln Ser Asp Gly Gln Ser Tyr
                                         285
                        280
Arg Gly Thr Ser Ser Thr Thr Ile Thr Gly Lys Lys Cys Gln Ser Trp
                    295
                                      300
Ala Ala Met Phe Pro His Arg His Ser Lys Thr Pro Glu Asn Phe Pro
                                  315
                 310
Asp Ala Gly Leu Glu Met Asn Tyr Cys Arg Asn Pro Asp Gly Asp Lys
             325
                              330
Gly Pro Trp Cys Tyr Thr Thr Asp Pro Ser Val Arg Trp Glu Tyr Cys
          340
                           345
Asn Leu Lys Arg Cys Ser Glu Thr Gly Gly
       355
     <210> 9
                          40 12 Feb
     <211> +552 ·
```

	< 2	21>			501												
	<2	22>	(1).	(5	52)												
cat His l	<4 act Thr	00> cat His	cag	gac Asp 5	ttt Phe	cag Gln	cca Pro	gtg Val	ctc Leu 10	cac His	ctg Leu	gtg Val	gca Ala	ctg Leu 15	aac Asn		48
acc Thr	ccc Pro	ctg Leu	tct Ser 20	gga Gly	ggc Gly	atg Met	cgt Arg	ggt Gly 25	atc Ile	cgt Arg	gga Gly	gca Ala	gat Asp 30	ttc Phe	cag Gln		96
tgc Cys	ttc Phe	cag Gln 35	caa Gln	gcc Ala	cga Arg	gcc Ala	gtg Val 40	Gly ggg	ctg Leu	tcg Ser	ggc Gly	acc Thr 45	ttc Phe	cgg Arg	gct Ala		.44
ttc Phe	ctg Leu 50	tcc Ser	tct Ser	agg Arg	ctg Leu	cag Gln 55	gat Asp	ctc Leu	tat Tyr	agc Ser	atc Ile 60	gtg Val	cgc Arg	cgt Arg	gct Ala	1	192
gac Asp 65	cgg Arg	Gly 999	tct Ser	gtg Val	ccc Pro 70	atc Ile	gtc Val	aac Asn	ctg Leu	aag Lys 75	gac Asp	gag Glu	gtg Val	cta Leu	ser 80		240
ccc Pro	agc Ser	tgg Trp	gac Asp	tcc Ser 85	ctg Leu	ttt Phe	tct Ser	ggc Gly	tcc Ser 90	cag Gln	ggt Gly	caa Gln	gtg Val	caa Gln 95	ccc Pro	2	288
GJA aaa	gcc Ala	cgc Arg	atc Ile 100	ttt Phe	tct Ser	ttt Phe	gac Asp	ggc Gly 105	aga Arg	gat Asp	gtc Val	ctg Leu	aga Arg 110	cac His	cca Pro	3	336
gcc Ala	tgg Trp	ccg Pro 115	cag Gln	aag Lys	agc Ser	gta Val	tgg Trp 120	His	ggc Gly	tcg Ser	gac Asp	ccc Pro 125	agt Ser	Gly aaa	cgg Arg	:	384
agg Arg	ctg Leu 130	atg Met	gag Glu	agt Ser	tac Tyr	tgt Cys 135	Glu	aca Thr	tgg Trp	cga Arg	act Thr 140	Glu	act Thr	act Thr	Gly 999	•	432
gct Ala 145	aca Thr	ggt	cag Gln	Ala	tcc Ser 150	Ser	Leu	Leu	Ser	Gly	agg Arg	Leu	Leu	gaa Glu	, cag. Gln 160	•	48.0
aaa Lys	gct Ala	gcg Ala	agc Ser	tgc Cys 165	His	aac Asn	ago Ser	tac Tyr	atc Ile 170	. Val	ctg Leu	tgc Cys	att Ile	gag Glu 175	aat Asn		528
agc Ser	ttc Phe	atg Met	acc Thr 180	Ser	ttc Phe	tcc Ser	aaa Lys	i i							·		552

<210> 10

<220>

<211> 184 <212> PRT <213> Mus musculus <400> 10 His Thr His Gln Asp Phe Gln Pro Val Leu His Leu Val Ala Leu Asn 10 Thr Pro Leu Ser Gly Gly Met Arg Gly Ile Arg Gly Ala Asp Phe Gln 25 Cys Phe Gln Gln Ala Arg Ala Val Gly Leu Ser Gly Thr Phe Arg Ala 40 Phe Leu Ser Ser Arg Leu Gln Asp Leu Tyr Ser Ile Val Arg Arg Ala 60 55 Asp Arg Gly Ser Val Pro Ile Val Asn Leu Lys Asp Glu Val Leu Ser 75 70 Pro Ser Trp Asp Ser Leu Phe Ser Gly Ser Gln Gly Gln Val Gln Pro 90 85 Gly Ala Arg Ile Phe Ser Phe Asp Gly Arg Asp Val Leu Arg His Pro 105 110 100 Ala Trp Pro Gln Lys Ser Val Trp His Gly Ser Asp Pro Ser Gly Arg 125 120 Arg Leu Met Glu Ser Tyr Cys Glu Thr Trp Arg Thr Glu Thr Thr Gly 140 135 Ala Thr Gly Gln Ala Ser Ser Leu Leu Ser Gly Arg Leu Leu Glu Gln 150 155 Lys Ala Ala Ser Cys His Asn Ser Tyr Ile Val Leu Cys Ile Glu Asn 170 165 Ser Phe Met Thr Ser Phe Ser Lys 180 <210> 11 <211> 1414 <212> DNA <213> Mus musculus <220> <221> CDS <222> (1) ... (1414) <400> 11 atg gac cat aag gaa gta atc ctt ctg ttt ctc ttg ctt ctg aaa cca 48 Met Asp His Lys Glu Val Ile Leu Leu Phe Leu Leu Leu Leu Lys Pro 10 1 gga caa ggg gac teg etg gat gge tac ata age aca caa ggg get tea 96 Gly Gln Gly Asp Ser Leu Asp Gly Tyr Ile Ser Thr Gln Gly Ala Ser 25 20 ctg ttc agt ctc acc aag aag cag ctc gca gca gga ggt gtc tcg gac Leu Phe Ser Leu Thr Lys Lys Gln Leu Ala Ala Gly Gly Val Ser Asp 40 35 tgt ttg gcc aaa tgt gaa ggg gaa aca gac ttt gtc tgc agg tca ttc 192 Cys Leu Ala Lys Cys Glu Gly Glu Thr Asp Phe Val Cys Arg Ser Phe

55

60

50

cag Gln 65	tac Tyr	cac His	agc Ser	aaa Lys	gag Glu 70	cag Gln	caa Gln	tgc Cys	gtg Val	atc Ile 75	atg Met	gcg Ala	gag Glu	aac Asn	agc Ser 80	240)
	act Thr															288	Ì
	gtg Val															336	;
gga Gly	acc Thr	atg Met 115	tcc Ser	agg Arg	aca Thr	aag Lys	agt Ser 120	ggt Gly	gtt Val	gcc Ala	tgt Cys	caa Gln 125	aag Lys	tgg Trp	ggt Gly	384	Ė
	acg Thr 130															432	?
	gga Gly															480)
	cct Pro															528	3
Asn	att Ile	Pro	Glu 180	Cys	Glu	Glu	Glu	Cys 185	Met	Tyr	Cys	Ser	Gly 190	Glu	Lys	57 <i>6</i>	
Tyr	gag Glu	Gly 195	Lys	Ile	Ser	Lys	Thr 200	Met	Ser	Gly	Leu	Asp 205	Cys	Gln	Ala	624	
Trp	gat Asp 210	Ser	Gln	Ser	Pro	His 215	Ala	His	Gly	Tyr	11e 220	Pro	Ala	Lys	Phe	672	
Pro 225	. agc. Ser	Lys	Asn	Leu	Lys 230	Met	Asn	Tyr	Cys	His 235	Asn	Pro	Asp	Gly	Glu 240	720	
Pro	agg Arg	Pro	Trp	Cys 245	Phe	Thr	Thr	Asp	Pro 250	Thr	Lys	Arg	Trp	Glu 255	Tyr	768	
tgt Cys	gac Asp	atc Ile	ccc Pro 260	cgc Arg	tgc Cys	aca Thr	aca Thr	ccc Pro 265	ccg Pro	ccc Pro	cca Pro	ccc Pro	agc Ser 270	cca Pro	acc Thr	816	
tac Tyr	caa Gln	tgt Cys	ctg Leu	aaa Lys	gga Gly	aga Arg	ggt Gly	gaa Glu	aat Asn	tac Tyr	cga Arg	ggg ggg	acc Thr	gtg Val	tct Ser	864	1

		275					280					285					
gtc Val	acc Thr 290	gtg Val	tct Ser	gly aaa	aaa Lys	acc Thr 295	tgt Cys	cag Gln	cgc Arg	tgg Trp	agt Ser 300	gag Glu	caa Gln	acc Thr	cct Pro		912
cat His 305	agg Arg	cac His	aac Asn	agg Arg	aca Thr 310	cca Pro	gaa Glu	aat Asn	ttc Phe	ccc Pro 315	tgc Cys	aaa Lys	aat Asn	ctg Leu	gaa Glu 320		960
gag Glu	aac Asn	tac Tyr	tgc Cys	cgg Arg 325	aac Asn	cca Pro	gat Asp	gga Gly	gaa Glu 330	act Thr	gct Ala	ccc Pro	tgg Trp	tgc Cys 335	tat Tyr	1	.008
acc Thr	act Thr	gac Asp	agc Ser 340	cag Gln	ctg Leu	agg Arg	tgg Trp	gag Glu 345	tac Tyr	tgt Cys	gag Glu	att Ile	cca Pro 350	tcc Ser	tgc Cys	1	.056
gag Glu	tcc Ser	tca Ser 355	gca Ala	tca Ser	cca Pro	gac Asp	cag Gln 360	tca Ser	gat Asp	tcc Ser	tca Ser	gtt Val 365	cca Pro	cca Pro	gag Glu	1	.104
gag Glu	caa Gln 370	aca Thr	cct Pro	gtg Val	gtc Val	cag Gln 375	gaa Glu	tgc Cys	tac Tyr	cag Gln	agc Ser 380	gat Asp	Gly 999	cag Gln	agc Ser	1	L152
tat Tyr 385	cgg Arg	ggt Gly	aca Thr	tcg Ser	tcc Ser 390	act Thr	acc Thr	atc Ile	aca Thr	999 Gly 395	aag Lys	aag Lys	tgc Cys	cag Gln	tcc Ser 400	1	1200
tgg Trp	gca Ala	gct Ala	atg Met	ttt Phe 405	cca Pro	cac His	agg Arg	cat His	tcg Ser 410	aag Lys	acc Thr	cca Pro	gag Glu	aac Asn 415	ttc Phe	3	1248
cca Pro	gat Asp	gct Ala	ggc Gly 420	ttg Leu	gag Glu	atg Met	aac Asn	tac Tyr 425	tgc Cys	agg Arg	aac Asn	ccg Pro	gat Asp 430	ggt Gly	gac Asp	:	1296
Lys	Gly	cct Pro 435	Trp	Cys	Tyr	Thr	Thr 440	Asp	Pro	Ser	· Val	Arg 445	Trp	Glu	Tyr	·	1344
tgc Cys	aac Asn 450	ctg Leu	aag Lys	cgg Arg	tgc Cys	tca Ser 455	Glu	aca Thr	gga Gly	ggg ggg	aat Asn 460	Ser	gac Asp	tac Tyr	aag Lys		1392
	Asp	gat Asp				taa *	C										1414

<210> 12

<2115 469 ·

<212> PRT

<213> Mus musculus

	< 4	100>	12												
Met 1	Asp	His	Lys	Glu 5	Val	Ile	Leu	Leu	Phe 10	Leu	Leu	Leu	Leu	Lys 15	Pro
Gly	Gln	Gly	Asp 20	Ser	Leu	Asp	Gly	Tyr 25	Ile	Ser	Thr	Gln	Gly 30	Ala	Ser
		Ser 35					40					45			
-	50	Ala				55					60				
65	-	His			70					75					80
-		Ser		85					90					95	
_		Tyr	100					105					110		
•		Met 115					120					125			
	130	Phe				135					140				
145	_	Leu			150					155					160
-		Trp		165					170					175	
		Pro	180					185					190		
-		Gly 195					200					205			
-	210	Ser				215					220				
225		Lys			230					235					240
		Pro		245					250					255	
_	_	Ile	260					265					270		
		Cys 275					280					285			
	290	Val				295					300				
305		His			310					315					320
		Tyr		325					330					335	
		Asp	340					345					350		
		Ser 355					360					365			
	370	Thr				375					380				
385	_	Gly			390					395					400
		Ala		405		25.5	A 7.	e:	419			'Æ	:Æ".	445	1
Pro	Asp	Ala	Gly 420	Leu	Glu	Met	Asn	Tyr 425	Cys	Arg	Asn	Pro	Asp 430	Gly	Asp

Lys	Gly	Pro 435	Trp	Cys	Tyr	Thr	Thr 440	Asp	Pro	Ser	Val	Arg 445	Trp	Glu	Tyr	
Cys	Asn 450	Leu	Lys	Arg	Cys	Ser 455	Glu	Thr	Gly	Gly	Asn 460	Ser	Asp	Tyr	Lys	
Asp 465	Asp	Asp	Asp	Lys												
	<2	210>	13													
	<2	211>	661													
		212>														
	<2	213>	Mus	muso	culus	3										
		20>														
		221>														
				(€	961)											
		.00>											_+_			48
													ctg Leu			40
Met 1	Asp	HIS	Lys	5	val	116	ьeu	пеп	10	Deu	Den	пеп	Deu	15	PIO	
1				,					10							
qqa	caa	qqq	gac	tcg	cta	gat	ctt	gac	tac	aag	gac	gac	gat	gac	aag	96
													Asp			
_			20					25					30			
													ctg			144
Leu	Ala		Thr	His	Gin	Asp		Gln	Pro	Val	Leu		Leu	vaı	Ala	
		35					40					45				
cta	aac	acc	ccc	cta	tct	gga	aac	atq	cat	aat	atc	cat	gga	qca	gat	192
													Gly			
	50					55	-		_	-	60					
													ggc			240
	Gln	Cys	Phe	GIn		Ala	Arg	Ala	Val		ьеп	Ser	Gly	Thr	Pne 80	
65					70					75					80	
caa	act	tta	cta	tac	tct	agg	cta	caq	gat	ctc	tat	agc	atc	ata	cqc	288
													Ile			
				85					90		-			95		
													gac			336
Arg	Ala	Asp		Gly	Ser	Val	Pro		Val	Asn	Leu	Lys	Asp	Glu	Val	
			100					105					110			
cta	tct	ccc	agc	taa	gac	tcc	cta	ttt	tct	aac	tcc	caq	ggt	caa	ata	384
													Gly			
		115		P	- r-		120			- 3		125	•			
													gtc			432
Gln		Gly	Ala	Arg	Ile		Ser	Phe	Asp	Gly		Asp	Val	Leu	Arg	
	130					135		_			140					
C 2 C	cca	acc	taa	cca	CaG	220	acc	at =	taa	Cac	aac	tcg	gac	כככ	agt	480
													Asp			
						4		_	-		-		-			

145					150					155					160	
Gly 999	cgg Arg	agg Arg	ctg Leu	atg Met 165	gag Glu	agt Ser	tac Tyr	tgt Cys	gag Glu 170	aca Thr	tgg Trp	cga Arg	act Thr	gaa Glu 175	act Thr	528
act Thr	Gly 999	gct Ala	aca Thr 180	ggt Gly	cag Gln	gcc Ala	tcc Ser	tcc Ser 185	ctg Leu	ctg Leu	tca Ser	ggc	agg Arg 190	ctc Leu	ctg Leu	576
gaa Glu	cag Gln	aaa Lys 195	gct Ala	gcg Ala	agc Ser	tgc Cys	cac His 200	aac Asn	agc Ser	tac Tyr	atc Ile	gtc Val 205	ctg Leu	tgc Cys	att Ile	624
	aat Asn 210									taa *	taa *	С				661
		220-	1.4													
		210> 211>														
		212>		mue	culus											
	٠.	2137	rids	ilius.	Jului	•										
	< Asp	400>		61	**- 1	T10	Lou	1 011	Dho	T ON	Lou	Len	Len	Luc	Pro	
met 1	Asp	HIS	гуѕ	5	vai	116	neu	neu	10	Бец	Беи	neu	<u> L</u> Cu	15	110	
Gly	Gln	Gly	Asp 20	Ser	Leu	Asp	Leu	Asp 25	Tyr	Lys	Asp	Asp	Asp	Asp	Lys	
Leu	Ala			His	Gln	Asp			Pro	Val	Leu	His 45		Val	Ala	
Leu	Asn	35 Thr	Pro	Leu	Ser		40 Gly	Met	Arg	Gly			Gly	Ala	Asp	
Dhe	50 Gln	Cvs	Phe	Gln	Gln	55 Ala	Ara	Ala	Val	Glv	60 Leu	Ser	Gly	Thr	Phe	
65		_			70					75					80	
Arg	Ala	Phe	Leu	Ser 85	Ser	Arg	Leu	Gln	Asp 90	Leu	Tyr	Ser	Ile	Val 95	Arg	
Arc	Ala	Asp	Arg		Ser	Val	Pro	Ile		Asn	Leu	Lys	Asp		Val	
_		_	100	_	_	_	.	105	9	03	0	03 =	110	01 =	Tro 1	
Leu	Ser	Pro 115	ser	Trp	Asp	ser	120	Pne	Ser	GIŸ	ser	125	GIY	GIII	Val	
Glr	Pro 130		Ala	Arg	.Ile	Phe 135		Phe	Asp	Gly	Arg 140	Asp	Val	Leu	Arg	
His	Pro	Ala	Trp	Pro	Gln		Ser	Val	Trp	His		Ser	Asp	Pro	Ser	
145					150	_	_			155	_	_	_,	~ 3	160	
Gly	Arg	Arg	Leu	Met 165	Glu	Ser	Tyr	Cys	Glu 170	Thr	Trp	Arg	Thr	Glu 175	Thr	
Thr	Gly	Ala	Thr 180		Gln	Ala	Ser	Ser 185	Leu	Leu	Ser	Gly	Arg 190	Leu	Leu	
Glu	Gln	_		Ala	Ser	Cys			Ser	Туr	Ile	Val 205		Cys	Ile	
Gli	Asn	195 Ser	Phe	Met	Thr	Ser	200 Phe	Ser	Lvs			203				
		eranti.				21.5			_				7	1		

<210> 15

	<	211> 212> 213>	DNA		culu	s										
	<	220> 221> 222>		(1747)										
_	gac		aag					_			ttg Leu		_			48
			-	_	_	-				-	aca Thr					96
_		_			_	_	_		_	_	gga Gly		_	_	_	144
_	_	_		_	-		_		_		gtc Val 60	_				192
_			_			-		_			atg Met		_			240
_							_	-	-	_	atc Ile			_	_	288
-			_		_	_	_				ggc Gly				_	336
											tgt Cys				_	384
					_						agt Ser 140				_	432
			_				_				gac Asp		_	-		480
			-				_	_	_	_	aga Arg		_		-	528
											tgc Cys					576

	180			1	85				190			
tat gag g Tyr Glu G	ggc aaa Bly Lys 195	atc tcc Ile Ser	Lys 1	acc at Thr Me 200	tg to et Se	t gga r Gly	ctt Leu	gac Asp 205	tgc Cys	cag Gln	gcc Ala	624
tgg gat t Trp Asp S 210	ct cag Ser Gln	agc cca Ser Pro	cat g His A 215	gct ca Ala Hi	at gg is Gl	a tac y Tyr	atc Ile 220	cct Pro	gcc Ala	aaa Lys	ttt Phe	672
cca agc a Pro Ser I 225												720
cca agg c Pro Arg F						o Thr						768
tgt gac a Cys Asp I				Chr P								816
tac caa t Tyr Gln C	gt ctg Cys Leu 275	aaa gga Lys Gly	Arg G	ggt g Sly G 280	aa aa lu As	t tac in Tyr	cga Arg	999 Gly 285	acc Thr	gtg Val	tct Ser	864
gtc acc g Val Thr V 290	gtg tct /al Ser	ggg aaa Gly Lys	acc t Thr C 295	gt ca Cys G	ag cg ln Ar	gc tgg gg Trp	agt Ser 300	gag Glu	caa Gln	acc Thr	cct Pro	912
cat agg o His Arg H 305												960
gag aac t Glu Asn T	tac tgc Tyr Cys	cgg aac Arg Asn 325	cca g Pro A	gat g Asp G	ga ga ly Gl 33	u Thr	gct Ala	ccc Pro	tgg Trp	tgc Cys 335	tat Tyr	1008
acc act o	gac agc Asp Ser 340	cag ctg Gln Leu	agg t Arg T	Trp G	ag ta lu Ty 45	c tgt r Cys	gag Glu	att Ile	cca Pro 350	tcc Ser	tgc Cys	1056
gag tcc t Glu Ser S	cca gca Ser Ala 355	tca cca Ser Pro	Asp (cag to Gln S 360	ca ga er As	t tcc sp Ser	tca Ser	gtt Val 365	cca Pro	cca Pro	gag Glu	1104
gag caa a Glu Gln 7 370	aca cct Thr Pro	gtg gga Val Gly	ggg a Gly 1 375	aat t Asn C	gc gg	gc ggt Ly Gly	gga Gly 380	tca Ser	ggt Gly	ggc Gly	gga Gly	1152
gat ctt g Asp Leu A 385	gac tac Asp Tyr	aag gac Lys Asp 390	gac q Asp A	gat g Asp A	sp Ly	ag ctt /s Leu 395	Ala	cat His	act Thr	cat His	cag Gln 140 marc	1200
gac ttt d	cag cca	gtg ctc	cac o	ctg g	tg go	ca ctg	aac	acc	ccc	ctg	tct	1248

Asp	Phe	Gln	Pro	Val 405	Leu	His	Leu	Val	Ala 410	Leu	Asn	Thr	Pro	Leu 415	Ser	
	ggc Gly															1296
	cga Arg															1344
	ctg Leu 450															1392
	ccc Pro								_							1440
	ctg Leu					_			-						_	1488
	tct Ser		-									_				1536
_	agc Ser	_				_	-		-							1584
_	tac Tyr 530	_				_										1632
	tcc Ser															1680
_	cac His		_			-	_	_				_		_		1728
	ttc Phe			taa *	taa *	C										1747
	<210> <211> <212> <213>		580 PRT	musc	culus	3										

28

<400> 16 Met Asp His Lys Glu Val Ile Leu Leu Phe Leu Leu Leu Leu Lys Pro

1				5					10					15	
	Gln	Gly	Asp 20		Leu	Asp	Gly	Tyr 25		Ser	Thr	Gln	Gly 30	Ala	Ser
		35	Leu				40					45			
-	50		Lys			55					60				
65	_		Ser		70					75					80
•			Ser	85					90					95	
			Leu 100					105					110		
_		115	Ser				120					125			
	130		Pro			135					140				
145	_		Glu		150					155					160
_			Cys	165					170					175	
			Glu 180					185					190		
-		195	Lys				200					205			
-	210		Gln			215					220				
225			Asn		230					235					240
			Trp	245					250					255	
-	_		Pro 260					265					270		
_		275	Leu				280					285			
	290		Ser			295					300				
305	_		Asn		310					315					320
			Cys	325					330					335	
			Ser, 340					345					350		
		355					360					365			Glu
	370		Pro			375					380				
385			Tyr		390					395					400
_			Pro	405					410					415	
8.	_		Arg 420			. :		¢425	4.1				4.3 G	asi J	١,
Ala	Arg	Ala 435	Val	Gly	Leu	Ser	Gly 440	Thr	Phe	Arg	Ala	Phe 445	Leu	ser	ser

WO 99/26480 PCT/US98/24950

```
Arg Leu Gln Asp Leu Tyr Ser Ile Val Arg Arg Ala Asp Arg Gly Ser
                        455
                                           460
Val Pro Ile Val Asn Leu Lys Asp Glu Val Leu Ser Pro Ser Trp Asp
                    470
                                         475
Ser Leu Phe Ser Gly Ser Gln Gly Gln Val Gln Pro Gly Ala Arg Ile
                                                         495
                485
                                    490
Phe Ser Phe Asp Gly Arg Asp Val Leu Arg His Pro Ala Trp Pro Gln
                                                     510
            500
                                505
Lys Ser Val Trp His Gly Ser Asp Pro Ser Gly Arg Arg Leu Met Glu
                                                525
                            520
        515
Ser Tyr Cys Glu Thr Trp Arg Thr Glu Thr Thr Gly Ala Thr Gly Gln
                                            540
                        535
Ala Ser Ser Leu Leu Ser Gly Arg Leu Leu Glu Gln Lys Ala Ala Ser
                    550
                                        555
Cys His Asn Ser Tyr Ile Val Leu Cys Ile Glu Asn Ser Phe Met Thr
                                    570
Ser Phe Ser Lys
            580
      <210> 17
      <211> 549
      <212> DNA
      <213> Homo sapiens
      <400> 17
cacagocaco gogaetteca geoggtgete cacetggttg egeteaacag ecceetgtea
                                                                        60
ggeggcatge ggggcatceg eggggcegac ttccagtgct tccagcagge gegggecgtg
                                                                       120
gggctggcgg gcaccttccg cgccttcctg tcctcgcgcc tgcaggacct gtacagcatc
                                                                       180
gtgcgccgtg ccgaccgcgc agccgtgccc atcgtcaacc tcaaggacga gctgctgttt
                                                                       240
                                                                       300
cccagctggg aggctctgtt ctcaggctct gagggtccgc tgaagcccgg ggcacgcatc
                                                                       360
ttctcctttg acggcaagga cgtcctgagg caccccacct ggccccagaa gagcgtgtgg
                                                                       420
catggctcgg accccaacgg gcgcaggctg accgagagct actgtgagac gtggcggacg
gaggeteect eggecaeggg ecaggeetee tegetgetgg ggggeagget eetggggeag
                                                                       480
                                                                       540
aqtqccqcqa gctgccatca cgcctacatc gtgctctgca ttgagaacag cttcatgact
                                                                       549
gcctccaag
      <210> 18
      <211> 183
      <212> PRT
      <213> Homo sapiens
      <400> 18
His Ser His Arg Asp Phe Gln Pro Val Leu His Leu Val Ala Leu Asn
1
                 5
Ser Pro Leu Ser Gly Gly Met Arg Gly Ile Arg Gly Ala Asp Phe Gln
                                25
Cys Phe Gln Gln Ala Arg Ala Val Gly Leu Ala Gly Thr Phe Arg Ala
                             40
Phe Leu Ser Ser Arg Leu Gln Asp Leu Tyr Ser Ile Val Arg Arg Ala
                        55
Asp Arg Ala Ala Val Pro Ile Val Asn Leu Lys Asp Glu Leu Leu Phe
                                        75
                    70
Pro Ser Trp Glu Ala LeumPhe Ser Gly Ser Glu Gly Pro Leu Lys Pro
                                     90
```

Gly Ala Arg Ile Phe Ser Phe Asp Gly Lys Asp Val Leu Arg His Pro

110

100

```
Thr Trp Pro Gln Lys Ser Val Trp His Gly Ser Asp Pro Asn Gly Arg
                                                  125
                             120
         115
 Arg Leu Thr Glu Ser Tyr Cys Glu Thr Trp Arg Thr Glu Ala Pro Ser
                         135
                                              140
 Ala Thr Gly Gln Ala Ser Ser Leu Leu Gly Gly Arg Leu Leu Gly Gln
                                          155
 Ser Ala Ala Ser Cys His His Ala Tyr Ile Val Leu Cys Ile Glu Asn
                 165
                                     170
 Ser Phe Met Thr Ala Ser Lys
       <210> 19
       <211> 5408
       <212> DNA
       <213> Homo sapiens
       <400> 19
                                                                        60
geogegtega egeggeggag gaggeageat eeegeggege tgaeggteet ggggagagea
                                                                       120
tggcgccgag gtgcccctgg ccatggccgc ggcggcggcg cctcctggac gtgctcgcgc
                                                                       180
ccctggtcct gctgctcggg gtccgcgcgg cctccgcgga gccagagcgc atcagcgagg
                                                                       240
aggtggggct gctgcagctc cttggggacc ccccgcccca gcaggtcacc cagacggatg
                                                                       300
accorgacyt cygyctyyce tacytettty yyccayatyc caacaytyyc caaytyycec
ggtaccactt ccccagcctc ttcttccgtg acttctcact gctgttccac atccggccag
                                                                       360
ccacagaggg cccaggggtg ctgttcgcca tcacggactc ggcgcaggcc atggtcttgc
                                                                       420
tgggcgtgaa gctctctggg gtgcaggacg ggcaccagga catctccctg ctctacacag
                                                                       480
aacctggtgc aggccagacc cacacagecg ccagetteeg geteeeegee ttegteggee
                                                                       540
agtggacaca cttagccctc agtgtggcag gtggctttgt ggccctctac gtggactgtg
                                                                       600
aggagttcca gagaatgccg cttgctcggt cctcacgggg cctggagctg gagcctggcg
                                                                       660
ccgggctctt cgtggctcag gcggggggag cggaccctga caagttccag ggggtgatcg
                                                                       720
ctgagctgaa ggtgcgcagg gacccccagg tgagccccat gcactgcctg gacgaggaag
                                                                       780
gcgatgactc agatggggca ttcggagact ctggcagcgg gctcggggac gcccgggagc
                                                                       840
ttctcaggga ggagacgggc gcggccctaa aacccaggct ccccgcgcca ccccccgtca
                                                                       900
ccacgccacc cttggctgga ggcagcagca cggaagattc cagaagtgaa gaagtcgagg
                                                                       960
agcagaccac ggtggcttcg ttaggagctc agacacttcc tggctcagat tctgtctcca
                                                                      1020
cgtgggacgg gagtgtccgg acccctgggg gccgcgtgaa agagggcggc ctgaaggggc
                                                                      1080
agaaagggga gccaggtgtt ccgggcccac ctggccgggc aggcccccca ggatccccat
                                                                      1140
gcctacctgg tcccccgggt ctcccgtgcc cagtgagtcc cctgggtcct gcaggcccag
                                                                      1200
cgttgcaaac tgtccccgga ccacaaggac ccccagggcc tccggggagg gacggcaccc
                                                                      1260
ctggaaggga cggcgagccg ggcgaccccg gtgaagacgg aaagccgggc gacaccgggc
                                                                      1320
cacaaggett ecctgggact ccaggggatg taggteecaa gggagacaag ggagaceetg
                                                                      1380
gggttggaga gagagggccc "ccaggacccc aagggcctcc agggccccca, ggaccctcct
                                                                      1440
tcaqacacga caagctgacc ttcattgaca tggagggatc tggctttggg ggcgatctgg
                                                                      1500
                                                                      1560
aggeeetgeg gggteetega ggetteeetg gaeeteeegg acceeeggt gteeeaggee
                                                                      1620
tgcccggcga gccaggccgc tttggggtga acagctccga cgtcccagga cccgccggcc
ttcctggtgt gcctgggcgc gagggtcccc ccgggtttcc tggcctcccg ggacccccag
                                                                      1680
gccctccggg aagagaggg cccccaggaa ggactgggca gaaaggcagc ctgggtgaag
                                                                      1740
                                                                      1800
caggegeece aggacataag gggageaagg gageeeeegg teetgetggt getegtgggg
                                                                      1860
agageggeet ggeaggagee eeeggacetg etggaceace aggeeeeeet gggeeeeetg
                                                                      1920
ggcccccagg accaggactc cccgctggat ttgatgacat ggaaggctcc ggggggccct
tetggtcaac agecegaage getgatggge cacagggace teceggeetg eegggactta
                                                                      1980
agggggatcc tggcgtgcct gggctgccgg gggcgaaggg agaagttgga gcagatggaa
Eccedggtt-ceceggecte*tettggddagaagaagtettge tgggccceaagaggddaaaaggaan 2189:00
gagacagagg cagccgggga gaaaagggag atccagggaa ggacggagtc gggcagccgg
                                                                      2160
geeteeetgg ceeeeegga eeeeegggae etgtggteta egtgteggag eaggaeggat
```

105

WO 99/26480 PCT/US98/24950

ccgtcctgag	cgtgccggga	cctgagggcc	ggccgggttt	cgcaggcttt	cccggacctg	2280
caggacccaa	gggcaacctg	ggctctaagg	gcgaacgagg	ctccccggga	cccaagggtg	2340
agaagggtga	accgggcagc	atcttcagcc	ccgacggcgg	tgccctgggc	cctgcccaga	2400
aaggagccaa	gggagagccg	ggcttccgag	gacccccggg	tccatacgga	cggccggggt	2460
acaagggaga	gattggcttt	cctggacggc	cgggtcgccc	cgggatgaac	ggattgaaag	2520
gagagaaagg	ggagccggga	gatgccagcc	ttggatttgg	catgagggga	atgcccggcc	2580
ccccaggacc	tccagggccc	ccaggccctc	cagggactcc	tgtttacgac	agcaatgtgt	2640
ttgctgagtc	cagccgcccc	gggcctccag	gattgccagg	gaatcagggc	cctccaggac	2700
ccaagggcgc	caaaggagaa	gtgggccccc	ccggaccacc	agggcagttt	ccgtttgact	2760
ttcttcagtt	ggaggctgaa	atgaaggggg	agaagggaga	ccgaggtgat	gcaggacaga	2820
aaggcgaaag	gggggagccc	gggggcggcg	gtttcttcgg	ctccagcctg	cccggccccc	2880
ccggcccccc	aggcccacgt	ggctaccctg	ggattccagg	tcccaaggga	gagagcatcc	2940
ggggccagcc	cggcccacct	ggacctcagg	gaccccccgg	catcggctac	gaggggcgcc	3000
agggccctcc	cggcccccca	ggccccccag	ggcccccttc	atttcctggc	cctcacaggc	3060
agactatcag	cgttcccggc	cctccgggcc	cccctgggcc	ccctgggccc	cctggaacca	3120
tagacacctc	ctcaggggtg	aggctctggg	ctacacgcca	ggccatgctg	ggccaggtgc	3180
acqaggttcc	cgagggctgg	ctcatcttcg	tggccgagca	ggaggagctc	tacgtccgcg	3240
tgcagaacgg	gttccggaag	gtccagctgg	aggcccggac	accactccca	cgagggacgg	3300
acaatgaagt	ggccgccttg	cagccccccg	tggtgcagct	gcacgacagc	aacccctacc	3360
cqcqqcqgga	gcacccccac	cccaccgcgc	ggccctggcg	ggcagatgac	atcctggcca	3420
qccccctcg	cctgcccgag	ccccagccct	accccggagc	cccgcaccac	agctcctacg	3480
tgcacctgcg	gccggcgcga	cccacaagcc	cacccgccca	cagccaccgc	gacttccagc	3540
cggtgctcca	cctggttgcg	ctcaacagcc	ccctgtcagg	cggcatgcgg	ggcatccgcg	3600
gggccgactt	ccagtgcttc	cagcaggcgc	gggccgtggg	gctggcgggc	accttccgcg	3660
ccttcctgtc	ctcgcgcctg	caggacctgt	acagcatcgt	gcgccgtgcc	gaccgcgcag	3720
ccqtqcccat	cgtcaacctc	aaggacgagc	tgctgtttcc	cagctgggag	gctctgttct	3780
caggetetga	gggtccgctg	aagcccgggg	cacgcatctt	ctcctttgac	ggcaaggacg	3840
tcctgaggca	cccacctgg	ccccagaaga	gcgtgtggca	tggctcggac	cccaacgggc	3900
gcaggctgac	cgagagctac	tgtgagacgt	ggcggacgga	ggctccctcg	gccacgggcc	3960
aggcctcctc	gctgctgggg	ggcaggctcc	tggggcagag	tgccgcgagc	tgccatcacg	4020
cctacatcgt	gctctgcatt	gagaacagct	tcatgactgc	ctccaagtag	ccaccgcctg	4080
gatgcggatg	gccggagagg	accggcggct	cggaggaagc	ccccaccgtg	ggcagggagc	4140
ggccggccag	cccctggccc	caggacctgg	ctgccatact	ttcctgtata	gttcacgttt	4200
catgtaatcc	tcaagaaata	aaaggaagcc	aaagagtgta	ttttttaaa	agtttaaaac	4260
agaagcctga	tgctgacatt	cacctgcccc	aactctcccc	tgacctgtga	gcccagctgg	4320
gtcaggcagg	gtgcagtatc	atgccctgtg	caacctcttg	gcctgatcag	accacggctc	4380
gatttctcca	ggatttcctg	ctttgggaag	ccgtgctcgc	cccagcaggt	gctgacttca	4440
tctcccacct	agcagcaccg	ttctgtgcac	aaaacccaga	cctgttagca	gacaggcccc	4500
gtgaggcaat	gggagctgag	gccacactca	gcacaaggcc	atctgggctc	ctccagggtg	4560
tgtgctcgcc	ctgcggtaga	tgggagggag	gctcaggtcc	ctggggctag	ggggagcccc	4620
ttctgctcag	ctctgggcca	ttctccacag	caaccccagg	ctgaagcagg	ttcccaagct	4680
cagaggcgca	ctgtgacccc	cagctccggc	ctgtcctcca	acaccaagca	cagcagcctg	4740
gggctggcct	cccaaatgag	ccatgagatg	atacatccaa	agcagacagc	tccaccctgg	4800
ccgagtccaa	gctgggagat	tcaagggacc	catgagttgg	ggtctggcag	cctcccatcc	4860
	tctcatgccc					4920
	gaacacggcc					4980
	ggcagtgagg					5040
	ggcagtcatt					5100
	aaccccgggg					5160
	ggtcctctgt					5220
	ctgtggaggg					5280
	acctgcctca					5340
	agcca a t toa					5400
aaaaaaa						5408

<210> 20 <211> 1336 <212> PRT <213> Homo sapiens


<400> 20 Met Ala Pro Arg Cys Pro Trp Pro Trp Pro Arg Arg Arg Leu Leu 10 Asp Val Leu Ala Pro Leu Val Leu Leu Gly Val Arg Ala Ala Ser 25 Ala Glu Pro Glu Arg Ile Ser Glu Glu Val Gly Leu Leu Gln Leu Leu 40 Gly Asp Pro Pro Pro Gln Gln Val Thr Gln Thr Asp Asp Pro Asp Val 55 60 Gly Leu Ala Tyr Val Phe Gly Pro Asp Ala Asn Ser Gly Gln Val Ala 75 70 Arg Tyr His Phe Pro Ser Leu Phe Phe Arg Asp Phe Ser Leu Leu Phe 85 90 His Ile Arg Pro Ala Thr Glu Gly Pro Gly Val Leu Phe Ala Ile Thr 105 110 100 Asp Ser Ala Gln Ala Met Val Leu Leu Gly Val Lys Leu Ser Gly Val 120 125 Gln Asp Gly His Gln Asp Ile Ser Leu Leu Tyr Thr Glu Pro Gly Ala 140 135 Gly Gln Thr His Thr Ala Ala Ser Phe Arg Leu Pro Ala Phe Val Gly 155 150 Gln Trp Thr His Leu Ala Leu Ser Val Ala Gly Gly Phe Val Ala Leu 165 170 Tyr Val Asp Cys Glu Glu Phe Gln Arg Met Pro Leu Ala Arg Ser Ser 190 185 180 Arg Gly Leu Glu Leu Glu Pro Gly Ala Gly Leu Phe Val Ala Gln Ala 200 205 195 Gly Gly Ala Asp Pro Asp Lys Phe Gln Gly Val Ile Ala Glu Leu Lys 215 220 Val Arg Arg Asp Pro Gln Val Ser Pro Met His Cys Leu Asp Glu Glu 230 235 Gly Asp Asp Ser Asp Gly Ala Phe Gly Asp Ser Gly Ser Gly Leu Gly 250 Asp Ala Arg Glu Leu Leu Arg Glu Glu Thr Gly Ala Ala Leu Lys Pro 265 Arg Leu, Pro Ala Pro Pro Pro Val Thr Thr Pro Pro Leu Ala Gly Gly 280 Ser Ser Thr Glu Asp Ser Arg Ser Glu Glu Val Glu Glu Gln Thr Thr 300 295 Val Ala Ser Leu Gly Ala Gln Thr Leu Pro Gly Ser Asp Ser Val Ser 315 310 Thr Trp Asp Gly Ser Val Arg Thr Pro Gly Gly Arg Val Lys Glu Gly 330 325 Gly Leu Lys Gly Gln Lys Gly Glu Pro Gly Val Pro Gly Pro Pro Gly 345 340 Arg Ala Gly Pro Pro Gly Ser Pro Cys Leu Pro Gly Pro Pro Gly Leu - 3000 sec • 30**95**0 sec • 3**95**0 sec • • Pro Cys Pro Val Ser Pro Leu Gly Pro Ala Gly Pro Ala Leu Gln Thr

	l Pro	Gly	Pro	Gln		Pro	Pro	Gly	Pro	Pro 395	Gly	Arg	Asp	Gly	Thr 400
38. Pr	5 o Gly	Arg	Asp	Gly	390 Glu	Pro	Gly	Asp	Pro		Glu	Asp	Gly	Lys	
	_	_,	-	405	01 -	a 1	D	D	410	m \	Desa	G1.	2	415	Clv.
	y Asp		420					425					430		
Pr	o Lys	Gly 435	Asp	Lys	Gly	Asp	Pro 440	Gly	Val	Gly	Glu	Arg 445	Gly	Pro	Pro
Gl [.]	y Pro		Gly	Pro	Pro	Gly		Pro	Gly	Pro	Ser		Arg	His	Asp
T.v	450 s Leu	Thr	Phe	Tle	Asp	455 Met	Glu	Glv	Ser	Glv	460 Phe	Glv	Glv	Asp	Leu
46	5				470					475					480
Gl	u Ala	Leu	Arg	Gly 485	Pro	Arg	Gly	Phe	Pro 490	Gly	Pro	Pro	Gly	Pro 495	Pro
Gl	y Val	Pro		Leu	Pro	Gly	Glu		Gly	Arg	Phe	Gly		Asn	Ser
Se	r Asp	Val	500 Pro	Gly	Pro	Ala	Gly	505 Leu	Pro	Gly	Val	Pro	510 Gly	Arg	Glu
al.	y Pro	515	Clu	Dho	Dro	Cly	520	Dro	cly	Pro	Pro	525 Gly	Pro	Pro	Gly
	530					535					540				
Ar	g Glu	Gly	Pro	Pro	Gly	Arg	Thr	Gly	Gln	Lys	Gly	Ser	Leu	Gly	Glu
54					550					555					560
Al	a Gly	Ala	Pro		His	Lys	Gly	Ser		Gly	Ala	Pro	Gly		Ala
_				565	_	_,	_	_ ,	570		_	01		575	61
Gl	y Ala	Arg	580	Glu	ser	GIY	Leu	585	GIY	Ala	Pro	GIY	590	Ala	GIY
Pro	o Pro	Gly 595	Pro	Pro	Gly	Pro	Pro 600	Gly	Pro	Pro	Gly	Pro 605	Gly	Leu	Pro
Ala	a Gly		Asp	Asp	Met	Glu 615	Gly	Ser	Gly	Gly	Pro 620	Phe	Trp	Ser	Thr
א ד	610 a Arg	Ca*	λΊэ	λεη	Glv		Gln	Glv	Pro	Dro		I.eu	Pro	Glv	Leu
62		SEL	AIG	тэр	630	FIO	0111	Gry	FIO	635	Cly	Deu		0 ±1	640
	s Gly	Asp	Pro	Glv		Pro	Glv	Leu	Pro		Ala	Lys	Gly	Glu	
•	_	-		645					650					655	
Gl	y Ala	Asp	Gly 660	He	Pro	GIÀ	Phe	Pro 665	Gly	Leu	Pro	GIY	Arg 670	GIU	GIY
Il	e Ala	Gly 675	Pro	Gln	Gly	Pro	Lys 680	Gly	Asp	Arg	Gly	Ser 685	Arg	Gly	Glu
Lag	s Gly		Dro	Glv	Lve	Asn		Val	Glv	Gln	Pro		Len	Pro	Glv
_	690	_				695					700				
	o Pro	Gly	Pro	Pro		Pro	Val	Val	Tyr		Ser	Glu	Gin	Asp	G1y 720
70	r Val	Ton	802	Va I	710 Pro	Gly	Dro	Glu	Glv	715	Dro	Glv	Dhe	Δla	
				725					730					735	
Pho	e Pro	Gly	Pro 740	Ala	Gly	Pro	Lys	Gly 745	Asn	Leu	Gly	Ser	Lys 750	Gly	Glu
Ar	g Gly	Ser 755	Pro	Gly	Pro	Lys	Gly 760	Glu	Lys	Gly	Glu	Pro 765	Gly	Ser	Ile
Ph	e Ser 770	Pro	Asp	Gly	Gly	Ala 775	Leu	Gly	Pro	Ala	Gln 780	Lys	Gly	Ala	Lys
Gl	y Glu	Pro	Gly	Phe	Arg	_	Pro	Pro	Gly			Gly	Arg	Pro	
78					790		_		_	795		_	_		800
Ty	r. L y s	Gły	GaLu	1.le 805	6 1y	Phe	Pro	Gly	Arg 810	Paco	Glya	Arg	Pro.	Gly 815	Metal · 🚓
As	n Gly	Leu	Lys	Gly	Glu	Lys	Gly	Glu	Pro	Gly	Asp	Ala	Ser	Leu	Gly

.: . . F.

	000				825					830		
Phe Gly Met	820	Met	Dro	Glv		Pro	Glv	Pro	Pro		Pro	Pro
Phe Gly Mec 835	AIG GIY	Mec	FIU	840	110	110	Cry	110	845			
Gly Pro Pro	Glv Thr	Pro	Val		Asp	Ser	Asn	Val	Phe	Ala	Glu	Ser
850			855	1	•			860				
Ser Arg Pro	Gly Pro	Pro	Gly	Leu	Pro	Gly	Asn	Gln	Gly	Pro	Pro	Gly
865		870					875					880
Pro Lys Gly	Ala Lys	Gly	Glu	Val	Gly	Pro	Pro	Gly	Pro	Pro	Gly	Gln
	885					890					895	
Phe Pro Phe	Asp Phe	Leu	Gln	Leu		Ala	Glu	Met	Lys		Glu	Lys
	900				905	_		_		910	_	
Gly Asp Arg	Gly Asp	Ala	Gly		Lys	Gly	Glu	Arg		GIu	Pro	GIA
915	-, -,		<u> </u>	920	.	D	~1	Dwa	925	C1	Dvo	Dwa
Gly Gly Gly	Phe Phe	GIY	935	ser	ьeu	Pro	GIY	940	PIO	GIY	PIO	PIO
930 Gly Pro Arg	Cl	Dro		T 1 0	Dro	Gly	Dro		Glv	Glu	Ser	Tle
945	GIY IYI	950	Gry	116	FIO	Gry	955	כעם	Cry	014		960
Arg Gly Gln	Pro Gly		Pro	Glv	Pro	Gln		Pro	Pro	Gly	Ile	
Arg Ory Cri	965			1		970	- 4			•	975	-
Tyr Glu Gly	Arg Glr	Gly	Pro	Pro	Gly	Pro	Pro	Gly	Pro	Pro	Gly	Pro
.,	980	-			985					990		
Pro Ser Phe	Pro Gly	Pro	His	Arg	Gln	Thr	Ile	Ser	Val	Pro	Gly	Pro
995				1000					1005			
Pro Gly Pro	Pro Gly	Pro	Pro	Gly	Pro	Pro	Gly			Gly	Ala	Ser
1010			1015			_		1020		~-3	~ 3	
Ser Gly Val	Arg Let			Thr	Arg	Gln			Leu	Gly	GIn	
1025	D 01.	1030		T	T1.	Dho	1035		Clu	Gln	Glu	104
His Glu Val			rp	Leu	iie	1050		Ата	GIU	GIII	1055	
Iou Tur Val	104		Δsn	Glv	Dhe			Val	Gln	Leu		
Leu Tyr Val	Arg Val		Asn	Gly		Arg		Val	Gln	Leu 1070	Glu	
_	Arg Val	Gln			1069	Arg	Lys			1070	Glu	Ala
Leu Tyr Val Arg Thr Pro	Arg Val 1060 Leu Pro	Gln			1069 Asp	Arg	Lys			1070 Ala	Glu	Ala
Arg Thr Pro	Arg Val 1060 Leu Pro 5	Gln Arg	Gly	Thr	1069 Asp	Arg S Asn	Lys Glu	Val	Ala 1089	1070 Ala 5	Glu) Leu	Ala Gln
Arg Thr Pro 107 Pro Pro Val 1090	Arg Val 1060 Leu Pro 5 Val Glr	Gln Arg	Gly His	Thr 1080 Asp	1069 Asp Ser	Arg Asn Asn	Lys Glu Pro	Val Tyr 1100	Ala 1089 Pro	1070 Ala S Arg	Glu Leu Arg	Ala Gln Glu
Arg Thr Pro 107 Pro Pro Val	Arg Val 1060 Leu Pro 5 Val Glr	Gln Arg	Gly His	Thr 1080 Asp	1069 Asp Ser	Arg Asn Asn	Lys Glu Pro Ala	Val Tyr 1100 Asp	Ala 1089 Pro	1070 Ala S Arg	Glu Leu Arg	Ala Gln Glu Ala
Arg Thr Pro 107 Pro Pro Val 1090 His Pro His 1105	Arg Val 1060 Leu Pro 5 Val Glr Pro Thr	Gln Arg Leu Ala 1110	Gly His 1099 Arg	Thr 1080 Asp S	1069 Asp Ser Trp	Arg Asn Asn Arg	Lys Glu Pro Ala	Val Tyr 1100 Asp	Ala 1089 Pro) Asp	1070 Ala S Arg	Glu Leu Arg Leu	Ala Gln Glu Ala 112
Arg Thr Pro 107 Pro Pro Val 1090 His Pro His	Arg Val 1060 Leu Pro 5 Val Glr Pro Thi	Gln Arg Leu Ala 1110	Gly His 1099 Arg	Thr 1080 Asp S	1069 Asp Ser Trp	Arg Asn Asn Arg Pro	Lys Glu Pro Ala 1115 Tyr	Val Tyr 1100 Asp	Ala 1089 Pro) Asp	1070 Ala S Arg	Glu Leu Arg Leu Pro	Ala Gln Glu Ala 112 His
Arg Thr Pro 107 Pro Pro Val 1090 His Pro His 1105 Ser Pro Pro	Arg Val 1060 Leu Pro 5 Val Glr Pro Thi Arg Leu	Gln Arg Leu Ala 1110 Pro	Gly His 1095 Arg O	Thr 1080 Asp Pro	Asp Ser Trp	Arg Asn Asn Arg Pro	Lys Glu Pro Ala 1115 Tyr	Val Tyr 1100 Asp Fro	Ala 1085 Pro) Asp	1070 Ala S Arg Ile	Leu Arg Leu Pro	Ala Glu Ala 112 His
Arg Thr Pro 107 Pro Pro Val 1090 His Pro His 1105	Arg Val 1060 Leu Pro 5 Val Glr Pro Thr Arg Leu 112 Tyr Val	Gln Arg Leu Ala 1110 Pro	Gly His 1095 Arg O	Thr 1080 Asp Pro	Asp Ser Trp Gln	Arg Asn Asn Arg Pro 1130 Ala	Lys Glu Pro Ala 1115 Tyr	Val Tyr 1100 Asp Fro	Ala 1085 Pro) Asp	1070 Ala Arg Ile Ala Ser	Leu Pro 1135	Ala Glu Ala 112 His
Arg Thr Pro 107 Pro Pro Val 1090 His Pro His 1105 Ser Pro Pro His Ser Ser	Arg Val 1060 Leu Pro 5 Val Glr Pro Thr Arg Leu Tyr Val 1140	Gln Arg Leu Ala 1110 Pro	Gly His 1095 Arg Glu Leu	Thr 1080 Asp Pro Pro	Asp Ser Trp Gln Pro	Arg Asn Asn Arg Pro 1130 Ala	Lys Glu Pro Ala 1115 Tyr Arg	Val Tyr 1100 Asp Pro	Ala 1089 Pro) Asp Gly	1070 Ala Arg Ile Ala Ser	Leu Pro 1135 Pro	Ala Glu Ala 112 His Pro
Arg Thr Pro 107 Pro Pro Val 1090 His Pro His 1105 Ser Pro Pro His Ser Ser Ala His Ser	Arg Val 1060 Leu Pro 5 Val Glr Pro Thr Arg Leu 112 Tyr Val 1140 His Arg	Gln Arg Leu Ala 1110 Pro	Gly His 1095 Arg Glu Leu	Thr 1080 Asp Pro Pro Arg	1069 Asp Ser Trp Gln Pro 1149 Pro	Arg Asn Asn Arg Pro 1130 Ala	Lys Glu Pro Ala 1115 Tyr Arg	Val Tyr 1100 Asp Pro	Ala 1089 Pro) Asp Gly	1070 Ala 5 Arg Ile Ala Ser 1150 Val	Leu Pro 1135 Pro	Ala Glu Ala 112 His Pro
Arg Thr Pro 107 Pro Pro Val 1090 His Pro His 1105 Ser Pro Pro His Ser Ser Ala His Ser	Arg Val 1060 Leu Pro 5 Val Glr Pro Thr Arg Leu 112 Tyr Val 1140 His Arg	Arg Leu Ala 1110 Pro	Gly His 1099 Arg Glu Leu	Thr 1080 Asp Pro Pro Arg Gln 1160	1069 Asp Ser Trp Gln Pro 1149 Pro	Arg Asn Asn Arg Pro 1130 Ala Val	Lys Glu Pro Ala 1115 Tyr Arg Leu	Val Tyr 1100 Asp Pro Pro	Ala 1085 Pro Asp Gly Thr Leu 1165	1070 Ala Arg Ile Ala Ser 1150 Val	Leu Arg Leu Pro 1135 Pro Ala	Ala Glu Ala 112 His Pro Leu
Arg Thr Pro 107 Pro Pro Val 1090 His Pro His 1105 Ser Pro Pro His Ser Ser Ala His Ser	Arg Val 1060 Leu Pro 5 Val Glr Pro Thr Arg Leu 112 Tyr Val 1140 His Arg	Arg Leu Ala 1110 Pro	Gly His 1099 Arg Glu Leu	Thr 1080 Asp Pro Pro Arg Gln 1160 Met	1069 Asp Ser Trp Gln Pro 1149 Pro	Arg Asn Asn Arg Pro 1130 Ala Val	Lys Glu Pro Ala 1115 Tyr Arg Leu	Val Tyr 1100 Asp Pro Pro	Ala 1085 Pro Asp Gly Thr Leu 1165 Gly	1070 Ala Arg Ile Ala Ser 1150 Val	Leu Arg Leu Pro 1135 Pro Ala	Ala Glu Ala 112 His Pro Leu
Arg Thr Pro 107 Pro Pro Val 1090 His Pro His 1105 Ser Pro Pro His Ser Ser Ala His Ser 115 Asn Ser Pro 1170	Arg Val 1060 Leu Pro 5 Val Glr Pro Thr Arg Leu 112 Tyr Val 1140 His Arg Leu Ser	Gln Arg Leu Ala 1110 Pro 5 His	Gly His 1095 Arg Glu Leu Phe Gly 1175	Thr 1080 Asp Pro Pro Arg Gln 1160 Met	1069 Asp Ser Trp Gln Pro 1149 Pro Arg	Arg Asn Asn Arg Pro 1130 Ala Val	Lys Glu Pro Ala 1115 Tyr Arg Leu Ile	Val Tyr 1100 Asp Pro Pro His Arg	Ala 1085 Pro) Asp Gly Thr Leu 1165 Gly	1070 Ala Arg Ile Ala Ser 1150 Val Ala	Leu Arg Leu Pro 1135 Pro Ala Asp	Ala Glu Ala 112 His Pro Leu Phe
Arg Thr Pro 107 Pro Pro Val 1090 His Pro His 1105 Ser Pro Pro His Ser Ser Ala His Ser 115 Asn Ser Pro 1170 Gln Cys Phe	Arg Val 1060 Leu Pro 5 Val Glr Pro Thr Arg Leu 112 Tyr Val 1140 His Arg Leu Ser Gln Glr	Gln Arg Leu Ala 1110 Pro 5 His Gly Asp Ala 1190	Gly His 1099 Arg Glu Leu Phe Gly 1179 Arg	Thr 1080 Asp Pro Pro Arg Gln 1160 Met	1069 Asp Ser Trp Gln Pro 1149 Pro Arg	Arg Asn Arg Pro 1130 Ala Colored Gly Gly	Lys Glu Pro Ala 1115 Tyr Arg Leu Ile Leu 1195	Val Tyr 1100 Asp Pro Pro His Arg 1180 Ala	Ala 1085 Pro) Asp Gly Thr Leu 1165 Gly)	1070 Ala Arg Ile Ala Ser 1150 Val Ala Thr	Leu Arg Leu Pro 1135 Pro Ala Asp	Ala Gln Glu Ala 112 His Pro Leu Phe Arg 120
Arg Thr Pro 107 Pro Pro Val 1090 His Pro His 1105 Ser Pro Pro His Ser Ser Ala His Ser 115 Asn Ser Pro 1170 Gln Cys Phe	Arg Val 1060 Leu Pro 5 Val Glr Pro Thr Arg Leu 112 Tyr Val 1140 His Arg Leu Ser Gln Glr	Gln Arg Leu Ala 1110 Pro 5 His Gly Asp Ala 1190	Gly His 1099 Arg Glu Leu Phe Gly 1179 Arg	Thr 1080 Asp Pro Pro Arg Gln 1160 Met	1069 Asp Ser Trp Gln Pro 1149 Pro Arg	Arg Asn Arg Pro 1130 Ala Coly Gly Leu	Lys Glu Pro Ala 1115 Tyr Arg Leu Ile Leu 1195 Tyr	Val Tyr 1100 Asp Pro Pro His Arg 1180 Ala	Ala 1085 Pro) Asp Gly Thr Leu 1165 Gly)	1070 Ala Arg Ile Ala Ser 1150 Val Ala Thr	Leu Arg Leu Pro 1135 Pro Ala Asp Phe Arg	Ala Glu Ala 112 His Pro Leu Phe Arg 120 Arg
Arg Thr Pro 107 Pro Pro Val 1090 His Pro His 1105 Ser Pro Pro His Ser Ser Ala His Ser 115 Asn Ser Pro 1170 Gln Cys Phe 1185 Ala Phe Leu	Arg Val 1060 Leu Pro 5 Val Glr Pro Thr Arg Leu 112 Tyr Val 1140 His Arg 5 Leu Ser Gln Glr Ser Ser	Gln Arg Leu Ala 1110 Pro 5 His Gly Asp Asp Asp Asp Asp Asp Asp Asp	Gly His 1095 Arg Glu Leu Phe Gly 1175 Arg	Thr 1080 Asp Pro Pro Arg Gln 1160 Met Ala	1069 Asp Ser Trp Gln Pro 1149 Pro Arg Val	Arg Asn Arg Pro 1130 Ala Gly Gly Leu 1210	Lys Glu Pro Ala 1115 Tyr Arg Leu Ile Leu 1195 Tyr	Val Tyr 1100 Asp Pro Pro His Arg 1180 Ala Ser	Ala 1085 Pro) Asp Gly Thr Leu 1165 Gly) Gly	IO70 Ala Arg Ile Ala Ser 1150 Val Thr Val	Leu Arg Leu Pro 1135 Pro Ala Asp Phe Arg 1215	Ala Gln Glu Ala 112 His Pro Leu Phe Arg 120 Arg
Arg Thr Pro 107 Pro Pro Val 1090 His Pro His 1105 Ser Pro Pro His Ser Ser Ala His Ser 115 Asn Ser Pro 1170 Gln Cys Phe	Arg Val 1060 Leu Pro 5 Val Glr Pro Thi Arg Leu 112 Tyr Val 1140 His Arg 5 Leu Sei Gln Glr Ser Sei 120 Ala Ala	Gln Arg Leu Ala 1110 Pro 5 His Gly Asp Asp Asp Asp Asp Asp Asp Asp	Gly His 1095 Arg Glu Leu Phe Gly 1175 Arg	Thr 1080 Asp Pro Pro Arg Gln 1160 Met Ala	1069 Asp Ser Trp Gln Pro 1149 Pro Arg Val Asp	Arg Asn Arg Pro 1130 Ala 5 Val Gly Gly Leu 1210 Asn	Lys Glu Pro Ala 1115 Tyr Arg Leu Ile Leu 1195 Tyr	Val Tyr 1100 Asp Pro Pro His Arg 1180 Ala Ser	Ala 1085 Pro) Asp Gly Thr Leu 1165 Gly) Gly	1070 Ala 5 Arg Ile Ala Ser 1150 Val 5 Ala Thr Val Glu	Leu Arg Leu Pro 1135 Pro Ala Asp Phe Arg 1215 Leu	Ala Gln Glu Ala 112 His Pro Leu Phe Arg 120 Arg
Arg Thr Pro 107 Pro Pro Val 1090 His Pro His 1105 Ser Pro Pro His Ser Ser Ala His Ser 115 Asn Ser Pro 1170 Gln Cys Phe 1185 Ala Phe Leu Ala Asp Arg	Arg Val 1060 Leu Pro 5 Val Glr Pro Thr Arg Leu 112 Tyr Val 1140 His Arg 5 Leu Ser Gln Glr Ser Ser Ala Ala 1220	Gln Arg Leu Ala 1110 Pro 5 His Gly Asp Asp Asp Asp Arg Arg Arg	Gly His 1095 Arg Glu Leu Phe Gly 1175 Arg Leu Pro	Thr 1080 Asp Pro Pro Arg Gln 1160 Met Ala Gln	Asp Ser Trp Gln Pro 1145 Pro Arg Val Asp Val 1225	Arg Asn Asn Arg Pro 1130 Ala 5 Val Gly Gly Leu 1210 Asn	Lys Glu Pro Ala 1115 Tyr Arg Leu Ile Leu 1195 Tyr	Val Tyr 1100 Asp Pro Pro His Arg 1180 Ala Ser	Ala 1085 Pro Asp Gly Thr Leu 1165 Gly Gly Ile	IO70 Ala Arg Ile Ala Ser 1150 Val Thr Val Glu 1230	Leu Arg Leu Pro 1135 Pro Ala Asp Phe Arg 1215 Leu	Ala Gln Glu Ala 112 His Pro Leu Phe Arg 120 Arg Leu
Arg Thr Pro 107 Pro Pro Val 1090 His Pro His 1105 Ser Pro Pro His Ser Ser Ala His Ser 115 Asn Ser Pro 1170 Gln Cys Phe 1185 Ala Phe Leu Ala Asp Arg	Arg Val 1060 Leu Pro 5 Val Glr Pro Thr Arg Leu 1120 Tyr Val 1140 His Arg 5 Leu Ser Gln Glr Ser Ser 120 Ala Ala 1220 Trp Glr	Gln Arg Leu Ala 1Pro 5 His Gly Asp Ala 1190 Arg S Arg	Gly His 1095 Arg Glu Leu Phe Gly 1175 Arg Leu Pro	Thr 1080 Asp Pro Pro Arg Gln 1160 Met Ala Gln Ile	1069 Asp Ser Trp Gln Pro 1149 Pro Arg Val Asp Val 1229 Ser	Arg Asn Asn Arg Pro 1130 Ala 5 Val Gly Gly Leu 1210 Asn	Lys Glu Pro Ala 1115 Tyr Arg Leu 119 Tyr Leu 1195 Tyr	Val Tyr 1100 Asp Pro Pro His Arg 1180 Ala Ser Lys	Ala 1085 Pro Asp Gly Thr Leu 1165 Gly Gly Ile Asp	1070 Ala 5 Arg Ile Ala Ser 1150 Val Thr Val Glu 1230 Pro	Leu Arg Leu Pro 1135 Pro Ala Asp Phe Arg 1215 Leu	Ala Gln Glu Ala 112 His Pro Leu Phe Arg 120 Arg Leu
Arg Thr Pro 107 Pro Pro Val 1090 His Pro His 1105 Ser Pro Pro His Ser Ser Ala His Ser 115 Asn Ser Pro 1170 Gln Cys Phe 1185 Ala Phe Leu Ala Asp Arg Phe Pro Ser	Arg Val 1060 Leu Pro 5 Val Glr Pro Thr Arg Leu 1120 His Arg 5 Leu Ser Gln Glr Ser Ser 1220 Trp Glr 5**	Gln Arg Leu Ala 1110 Pro 5 His Gly Asp Asp Ala 1190 Arg N Arg N Arg N Arg Arg Arg Arg Arg Arg Arg Arg Arg	Gly His 1095 Arg Glu Leu Phe Gly 1175 Arg Leu Pro	Thr 1080 Asp Pro Pro Arg Gln 1160 Met Gln Ile	Asp Ser Trp Gln Pro 1145 Pro Arg Val Asp Val 1225 Ser	Arg Asn Arg Pro 1130 Ala Coly Gly Leu 1210 Asn Coly	Lys Glu Pro Ala 1115 Tyr Arg Leu 1195 Tyr Leu Ser	Val Tyr 1100 Asp Pro Pro His Arg 1180 Ala Ser Lys Glu	Ala 1085 Pro Asp Gly Thr Leu 1165 Gly Ile Asp	IOTO Ala Arg Ile Ala Ser 1150 Val Thr Val Glu 1230 Pro	Leu Arg Leu Pro 1135 Pro Ala Asp Phe Arg 1215 Leu Leu	Ala Gln Glu Ala 112 His Pro Leu Phe Arg 120 Arg Leu Lys
Arg Thr Pro 107 Pro Pro Val 1090 His Pro His 1105 Ser Pro Pro His Ser Ser Ala His Ser 115 Asn Ser Pro 1170 Gln Cys Phe 1185 Ala Phe Leu Ala Asp Arg	Arg Val 1060 Leu Pro 5 Val Glr Pro Thr Arg Leu 1120 His Arg 5 Leu Ser Gln Glr Ser Ser 1220 Trp Glr 5**	Gln Arg Leu Ala 1110 Pro 5 His Gly Asp Asp Ala 1190 Arg N Arg N Arg N Arg Arg Arg Arg Arg Arg Arg Arg Arg	Gly His 1095 Arg Glu Leu Phe Gly 1175 Arg Leu Pro	Thr 1080 Asp Pro Pro Arg Gln 1160 Met Ala Gln Ile Phe 12:0	Asp Ser Trp Gln Pro 1145 Pro Arg Val Asp Val 1225 Ser	Arg Asn Arg Pro 1130 Ala Coly Gly Leu 1210 Asn Coly	Lys Glu Pro Ala 1115 Tyr Arg Leu 1195 Tyr Leu Ser	Val Tyr 1100 Asp Pro Pro His Arg 1180 Ala Ser Lys Glu	Ala 1085 Pro Asp Gly Thr Leu 1165 Gly Gly Ile Asp Gly 1245 Val	IOTO Ala Arg Ile Ala Ser 1150 Val Thr Val Glu 1230 Pro	Leu Arg Leu Pro 1135 Pro Ala Asp Phe Arg 1215 Leu Leu	Ala Gln Glu Ala 112 His Pro Leu Phe Arg 120 Arg Leu Lys

WO 99/26480 PCT/US98/24950



International application No. PCT/US98/24950


IPC(6) :A01N US CL : 424/	ICATION OF SUBJECT MATTER N 63/00, 43/04; C12N 15/00; C07H 21/02 /93.1; 435/320.1; 514/44; 536/23.1		
	emational Patent Classification (IPC) or to both	national classification and IPC	
	SEARCHED CHARLES OF THE CONTROL OF T		
1	entation searched (classification system follow	ed by classification symbols)	
U.S. : 424/9	93.1; 435/320.1; 514/44; 536/23.1		
Documentation se	arched other than minimum documentation to the	e extent that such documents are included	in the fields searched
Electronic data ba	ase consulted during the international search (n	ame of data base and where practicable	search terms used)
Please See Extra	- ,		,
C. DOCUME	NTS CONSIDERED TO BE RELEVANT		
Category* C	Citation of document, with indication, where as	ppropriate, of the relevant passages	Relevant to claim No.
	5,792,845 A (O'REILLY et al.) . 4, lines 32-68, col. 5, lines 1-2,	1-30, 33	
СО	O 97/23500 A1 (THE CHILDR PRPORATION) 03 July 1997 (03. ge 42, lines 1-27.	4	
) 98/49321 A2 (RHONE-POULE 98 (05.11.98), page 44, 6-11, 25-33	1, 5, 18, 20, 31	
Y,P	0 (05.11.70), page 44, 0-11, 25-35	, page 43, imes 12-13, 29-33.	2-4, 6-17, 19, 21- 30, 32, 33
			,
X Further doc	suments are listed in the continuation of Box C	See patent family annex.	
-	egories of cited documents:	"T" later document published after the inte date and not in conflict with the appl	
A document d to be of par	defining the general state of the art which is not considered uticular relevance	the principle or theory underlying the	
	ument published on or after the international filing date	"X" document of particular relevance; the considered novel or cannot be conside	e claimed invention cannot be red to involve an inventive step
cited to est	which may throw doubts on priority claim(s) or which is tablish the publication date of another citation or other	when the document is taken alone "Y" document of particular relevance: the	
	son (as specified) referring to an oral disclosure, use, exhibition or other	"Y" document of particular relevance; the considered to involve an inventive	step when the document is
P document p	published prior to the international filing date but later than	being obvious to a person skilled in t	he art
	completion of the international search	Date of mailing of the international sea	
04 FEBRUARY		08 MAR 19	•
Name and mailing Commissioner of P Box PCT Washington, D.C.	address of the ISA/US Patents and Trademarks 20231	Authorized office A	7 Jen / 9
•	703) 305-3230	Telephone No. (703) 308-0196	(

INTERNATIONAL SEARCH REPORT

International application No. PCT/US98/24950

C (Continua	ation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant	passages	Relevant to claim No.
Y	WO 97/15666 A (THE CHILDREN'S MEDICAL CENT) CORPORATION) 01 May 1997 (01.05.97), page 20, line page 21, page 22, lines 1-16. page 59, lines 5-35, page 66 61, 1-6.	1-30, 33	
Y	TANAKA, T. et al. Retroviral and adenoviral mediated transduction of angiostatin cDNA inhibits angiogenesis at growth. Proceedings of the American Association for Car Research. March 1997 (03.97). Vol 38. page 264.	1-33	
Y	WO 96/35774 A2 (THE CHILDREN'S MEDICAL CENT CORPORATION) 14 November 1996 (14.11.96), page lin pages 22-25, page 26, lines 1-33. pages 144-148.	1-30, 33	
Y	WO 97/41824 A2(ABBOTT LABORATORIES) 13 Nove 1997 (13.11.97), page 5, lines 13-38, page 6, 1-18, page 6 15-38, pages 61-62, page 63, lines 1-33.	1-30, 33	
Y	WO 95/29242 A1 (THE CHILDREN; S MEDICAL CENT CORPORATION) 02 November 1995 (02.11.95), page 2: 19-35, pages 22-27, page 28, lines 1-4. page 87, lines 4-3 88, page 89, lines 1-14.	1-30, 33	
Y	O'REILLY, et al. Angiostatin induces and sustains dormal human primary tumors in mice. Nature Medicine, June 19 (06.96), Vol. 2, No. 6, pages 689-692, especially pages 68	1-30, 33	
Y	O'REILLY, et al. Endostatin: An Endogenous Inhibitor of Angiogenesis and Tumor Growth. Cell, 24 January 1997 (24.01.97), Vol. 88, pages 277-285, especially pages 279-		1-30, 33
,	•• :		¢. :

Form PCT/ISA/210 (continuation of second sheet)(July 1992)*





PCT/US98/24950

B. FIELDS SEARCHED Electronic data bases consulted (Name of data base and where practicable terms used):										
APS, STN, WPIDS, MEDLINE, CAPLUS, BIOSIS, EMBASE search terms: angiostatin, plasminogen, endostatin, colagen(w) XVIII, inhibit?(5a)tumor(5a)growth, tumor(5a)regress?, diabet?(p)retinopathy, plasmid, viral(5a) vector.										
	ļ									
and the second of the second o										
\cdot	-									

Form PCT/ISA/210 (extra sheet)(July 1992)*

This Page Blank (uspto)

This Page Blank (uspto)